

***R*-matrix calculation of the continuum states of carbon monoxide**

K Chakrabarti¹ and Jonathan Tennyson²

¹ Department of Mathematics, Scottish Church College, 1 and 3 Urquhart Sq., Kolkata 700 006, India

² Department of Physics and Astronomy, University College London, Gower St., London WC1E 6BT, UK

Received 2 March 2007, in final form 27 March 2007

Published 24 May 2007

Online at stacks.iop.org/JPhysB/40/2135

Abstract

Calculations on electron collisions with CO⁺ molecular ion are presented as a function of electron energy and CO⁺ geometry. Resonance positions and widths are obtained for the low-lying Feshbach resonances in the system. Plots of resonance curves suggest that low-energy dissociative recombination can occur via resonances of ¹Π, ¹Δ, ³Σ⁺ and ³Π symmetries. Electron impact electronic excitation is considered both for excitation to the first excited A ²Π state and to yield a dissociation cross section. The latter calculations suggest that the lower of two experimental measurements of this process is likely to be correct. Finally results are presented for bound states in the continuum for both ¹Σ⁻ and ³Σ⁻ symmetries.

(Some figures in this article are in colour only in the electronic version)

1. Introduction

Carbon monoxide is the second most abundant molecule in the Universe and its ion, CO⁺, has been observed in a variety of astronomical locations (Erickson *et al* 1981, Storzer *et al* 1995, Fuente and Martin Pintado 1997, Fox and Hac 1999). Given that such ions occur in plasmas, collisions with low-energy electrons are likely to be important. In particular, low-energy electron collisions can lead to both (de-)excitation of the ion and to its destruction.

At low collision energies dissociative recombination (DR) is the major route to destruction for molecular ions. Low-energy DR rates for CO⁺ have been measured both in merged beams (Mitchell and Hus 1985) and storage rings (Rosen *et al* 1998). Models of dissociative recombination require both accurate resonance curves and associated widths, and a detailed treatment of the nuclear motion problem (Schneider *et al* 2000, Florescu-Mitchell and Mitchel 2006). Here we provide the curves that could provide a starting point for such a treatment.

Besides DR studies, there are few studies of electron–CO⁺ collisions. A recent measurement focused on electron impact ionization (Lecointre *et al* 2006), and the two cited

DR studies (Mitchell and Hus 1985, Rosen *et al* 1998) also presented results for electron impact dissociation which were not in good agreement with each other. Rather surprisingly, given its importance, we know of no previous theoretical study of continuum states of CO⁺.

In an earlier study (Chakrabarti and Tennyson 2006), we used the *R*-matrix method to present a comprehensive study of the Rydberg states of the CO. In this work we consider the low-lying continuum of the CO system placing particular emphasis on those resonances that are likely to be important for dissociative recombination. We also consider processes involving electron impact excitation. The following section outlines the general method and details the calculations. Results are presented in section 3 and the final section presents our conclusions.

2. Calculations

2.1. Method

The details of the calculations are already presented in our earlier work (Chakrabarti and Tennyson 2006) so we present here only the essentials. The *R*-matrix method divides configuration space into two regions (Burke and Berrington 1993). The inner region is defined by a sphere—here of radius $10a_0$ —centred at the molecular centre of mass. This sphere encloses the entire *N*-electron target wavefunction, where for present purposes the target' means CO⁺. In this inner region, the wavefunction of the (*N* + 1)-electron system, CO, is given by

$$\Psi_k = \mathcal{A} \sum_{i,j} a_{i,j,k} \Phi_i(1, \dots, N) F_{i,j}(N+1) + \sum_i b_{i,k} \chi_i(1, \dots, N+1), \quad (1)$$

where \mathcal{A} is the anti-symmetrization operator, $F_{i,j}$ are continuum orbitals and χ_i are two-centre L^2 functions constructed from *N*-electron 'target' orbitals. Also Φ_i is the wavefunction of the *i*th target state. Electron-correlation effects are included in these target wavefunctions via configuration interaction (CI) expansions (Tennyson 1996a). The variational coefficients $a_{i,j,k}$ and $b_{i,k}$ are determined by diagonalizing the Hamiltonian matrix. For diatomic calculations, the inner region code is based on the Alchemy (McLean 1971) quantum chemistry package.

2.2. CO⁺ target

Calculations on both CO⁺ and CO were performed for ten internuclear separations in the range 1.5–3.5 a_0 . The CO Slater-type orbitals of Kirby-Docken and Liu (1977) were used to build a CO⁺ molecular basis of 46 molecular orbitals (24 σ , 14 π , 6 δ , 2 ϕ). An initial set of molecular orbitals was obtained by performing self-consistent field (SCF) calculations on the two lowest states of CO⁺, X ² Σ^+ and A ² Π . These SCF molecular orbitals were then used in a configuration interaction (CI) calculation. In all calculations the 1 σ and 2 σ molecular orbitals were frozen and the CI calculation used the following configurations:

$$\begin{aligned} &(3\sigma, 4\sigma, 5\sigma, 1\pi)^9, \\ &(3\sigma, 4\sigma, 5\sigma, 1\pi)^8 (6\sigma - 24\sigma, 2\pi - 14\pi, 1\delta - 6\delta)^1, \\ &(3\sigma, 4\sigma, 5\sigma, 1\pi)^7, (6\sigma - 24\sigma, 2\pi - 14\pi, 1\delta - 6\delta)^2. \end{aligned}$$

These correspond, respectively, to a complete active space (CAS) spanning valence orbitals, single excitations from this CAS and double excitations. Two sets of pseudo-natural orbitals (NOs) were obtained from CI calculation on the lowest CO⁺ states with ² Σ^+ and ² Π symmetry described above. Following Tennyson (1996b), all σ and the 1 π orbital in the target

state wavefunction were represented by ${}^2\Sigma^+$ NOs, and the remaining π and all δ orbitals were represented by the ${}^2\Pi$ NOs.

Our model uses the NOs constructed as above, 4 electrons frozen the $1\sigma^2$ and $2\sigma^2$ orbitals corresponding to the O and C 1s electrons respectively, and a $(3\sigma, 4\sigma, 5\sigma, 6\sigma, 1\pi, 2\pi)^9$ CAS-CI (note that in our earlier work the CAS used was wrongly quoted as $(3\sigma, 4\sigma, 5\sigma, 1\pi)^9$). The CO^+ excitations were compared by Chakrabarti and Tennyson (2006) with the experimental results of Huber and Herzberg (1979) and the CCSDT calculation of Okada and Iwata (2000) and were generally found to be in good agreement with our work. Calculated potential energy curves for the 14 target states used in the close coupling expansion can also be found in our previous paper.

2.3. The CO model

Our calculations used 14 CO^+ natural orbitals ($8\sigma, 4\pi, 2\delta$). These orbitals were augmented by continuum orbitals $F_{i,j}$, expressed as a truncated partial wave expansion around the centre of mass. Partial waves with $l \leq 6$ and $m \leq 2$ were retained in the expansion. The radial parts of the continuum functions were generated as numerical solutions of an isotropic Coulomb potential. Those solutions with an energy below 10 Ryd were retained. A Buttle (1967) correction was used to correct for the effect of this truncation or, alternatively, the fixed boundary condition used to generate the functions. To correct for linear dependence effects two σ and one π orbitals were removed using Lagrange orthogonalization (Tennyson *et al* 1987). The resulting $62\sigma, 52\pi, 42\delta$ functions were Schmidt orthogonalized to the target NOs.

Calculations were performed using the $(3\sigma-6\sigma, 1\pi, 2\pi)^9$ CAS target wavefunction for the CO^+ states, up to 14 of which were used in the close-coupling expansion (1). Further details of the calculation, including a tabulation of the target states and energies involved, can be found in our previous study (Chakrabarti and Tennyson 2006). Calculations were performed for the lowest four symmetries of both singlet and triplet symmetry, meaning that the resonances we consider are of total symmetry ${}^1\Sigma^+, {}^1\Pi, {}^1\Delta, {}^1\Sigma^-, {}^3\Sigma^+, {}^3\Pi, {}^3\Delta$ and ${}^3\Sigma^-$. Resonance series belonging to angular momentum symmetries Φ or higher will be both of higher energy and have narrower resonance meaning that these resonances are of less physical significance. As we only consider electronic excitation here, no attempt was made to top up the calculation for neglected symmetries. Experience shows that these higher symmetries, while important for elastic scattering, do not contribute significantly to electron impact electronic excitation cross sections (Gorfinkiel *et al* 2002).

2.4. Resonances

The inner region solutions, equation (1), were used to construct R -matrices on the boundary. In the outer region, in addition to the Coulomb potential, the potential was given by the diagonal and off-diagonal dipole and quadrupole moments of the CO^+ target states. The R -matrices were propagated (Morgan 1984) in this potential until the wavefunction could be matched with exponentially decreasing functions obtained from a Gailitis expansion (Noble and Nesbet 1984). Tests showed that propagating the R -matrices to $100a_0$ gave stable results and this value was used for all calculations presented here. Not all target states included in the inner region close-coupling expansion were included in the outer region calculation as those which are strongly closed were found to make negligible difference in the calculation.

Resonances were detected and fitted to a Breit–Wigner profile to obtain their energy (E_r) and width (Γ) using the RESON program (Tennyson and Noble 1985) with an energy grid

0.25×10^{-3} Ryd. The magnitudes of the complex quantum defects $\mu = \alpha + i\beta$ were obtained using the relations

$$E_r = E_t - \frac{1}{\nu^2}, \quad \Gamma = \frac{2\beta}{\nu^3} \quad (2)$$

where the effective quantum number, ν , equals $n - \alpha$ and E_t is the energy of the threshold to which the Rydberg series converges.

Although this work is confined to energies higher than the CO ionization limit some of the states we find are bound within our model. These states are of Σ^- electronic symmetry which cannot couple to the $X^2\Sigma^+$ ground state of CO^+ by loss of a single electron. Energies of these states thus appear as bound states associated with the Rydberg series converging on the $A^2\Pi$ first excited state of CO^+ . These bound states were found using program BOUND which employs the searching algorithm of Sarpal *et al* (1991) with an improved nonlinear, quantum defect based grid (Rabadan and Tennyson 1996).

3. Results

3.1. Resonances

There is clearly a considerable quantity of data available for the resonances from these calculations so all we can do here is present a summary. Further information or numerical values for data presented in the figures can be obtained from the authors.

Tables 1 and 2 present data on the resonances obtained for the CO equilibrium bondlength of $2.132a_0$ for singlet and triplet total symmetries respectively. In each case we tabulate values for resonance series converging on the lowest two electronically excited for which coupling is allowed. For the triplets these correspond to resonances associated with the $A^2\Pi$ and a $4\Sigma^+$ states respectively, which are indeed the lowest two excited states of CO^+ in the equilibrium region. However, the singlet resonances cannot couple to quartet target states so for the singlets the first two resonance series converge to the $A^2\Pi$ and $B^2\Sigma^+$ states.

Tables 1 and 2 also give the effective quantum number, ν . This parameter is useful for identifying both resonances which form a series and possible intruder states, i.e., resonances which are actually associated with a higher-lying excited state of the ion. Thus, for example, $1\Sigma^+$ symmetry resonances show a clear series of resonances with the real part of the quantum defect, α , equal to about 0.6. Conversely the resonance at 0.1386 Ryd has a quantum defect of 0.4, unlike any other resonance detected in this series, and is almost certainly an intruder or interloper (McLaughlin *et al* 2001) state. Such states can also be detected by analysing their behaviour as a function of internuclear separation, R . This can be done by either plotting the variation of the quantum defects with R (Rabadán and Tennyson 1997), or just the resonance curves which is particularly effective if the ionic curves they are associated with behave differently with R . Figures 1 and 2 show the behaviour of the low-lying two resonances series as a function of R for the symmetries for which we find low-lying resonances.

The most important resonances for dissociative recombination (DR) studies are those low-lying ones which lead to dissociation by crossing the target ion ground state potential energy curve at energies low enough to be energetically accessible. For states which become bound, i.e. cross the $\text{CO}^+ X^2\Sigma^+$ curve, within the range of bondlengths considered we have also attempted to determine the location of the corresponding bound states region of the curve using our bound state detecting program BOUND (Sarpal *et al* 1991). This results, as can be seen from the figure, in curves that link the quasi-bound and bound states, but that these curves are often far from smooth. This lack of smoothness is not due to noise in our calculations but is caused by couplings to the (infinite) Rydberg series converging to the $\text{CO}^+ X^2\Sigma^+$

Table 1. Resonance positions and widths (in Ryd), and effective quantum numbers (ν) at $R = 2.132a_0$ for singlet total symmetries of the CO molecule below the first two CO^+ excited states. Numbers within brackets indicate power of 10.

Below $^2\Pi$ state			Below $^2\Delta$ state		
Position	Width	ν	Position	Width	ν
$^1\Sigma^+$					
0.1090	0.2068 (-02)	2.3638	0.3172	0.3140 (-02)	2.4091
0.1386	0.2722 (-01)	2.5870	0.3712	0.1104 (-02)	2.9078
0.1992	0.4533 (-02)	3.3564	0.3852	0.4259 (-02)	3.0964
0.2006	0.5409 (-02)	3.3833	0.3873	0.4953 (-02)	3.1288
0.2254	0.1355 (-04)	3.9970	0.4059	0.9631 (-03)	3.4600
0.2317	0.1193 (-02)	4.2135	0.4237	0.4112 (-03)	3.8988
0.2368	0.7869 (-03)	4.4206	0.4269	0.4054 (-04)	3.9967
0.2477	0.6932 (-04)	4.9834	0.4308	0.1285 (-02)	4.1292
0.2540	0.2918 (-03)	5.4211	0.4393	0.4684 (-03)	4.4664
0.2580	0.8255 (-03)	5.7791	0.4477	0.1802 (-03)	4.8921
0.2637	0.1455 (-03)	6.4203	0.4515	0.6514 (-03)	5.1352
0.2654	0.9502 (-03)	6.6568	0.4560	0.2608 (-03)	5.4718
0.2698	0.9090 (-04)	7.4192	0.4606	0.9713 (-04)	5.8889
0.2706	0.7386 (-03)	7.5934			
$^1\Pi$					
0.0376	0.2137 (-03)	1.9983	0.3076	0.6750 (-03)	2.3451
0.1155	0.7100 (-03)	2.4076	0.3852	0.6825 (-03)	3.0973
0.1696	0.4020 (-03)	2.9068	0.4008	0.1995 (-03)	3.3594
0.1772	0.7000 (-03)	3.0044	0.4216	0.3425 (-02)	3.8404
0.1847	0.1181 (-03)	3.1113	0.4307	0.5004 (-03)	4.1266
0.2035	0.2352 (-04)	3.4398	0.4370	0.7934 (-04)	4.3660
0.2219	0.1150 (-03)	3.8897	0.4514	0.1994 (-03)	5.1235
0.2375	0.7976 (-05)	4.4508	0.4547	0.4684 (-04)	5.3672
0.2500	0.1713 (-04)	5.1335	0.4648	0.3093 (-04)	6.3676
0.2591	0.2736 (-04)	5.8801			
$^1\Delta$					
0.1110	0.5150 (-03)	2.3769	0.3402	0.9520 (-03)	2.5886
0.1969	0.3472 (-03)	3.3136	0.3546	0.1335 (-01)	2.7227
0.2023	0.1848 (-04)	3.4168	0.3782	0.8065 (-03)	2.9985
			0.4264	0.1065 (-03)	3.9810
			0.4495	0.9671 (-05)	5.0005
			0.4689	0.9671 (-05)	6.9708

ionization limit which couple with dissociative intruder states. Our curves therefore represent (approximate) diabats for these intruders.

Figures 1 and 2 show that the lowest resonance with $^1\Pi$, $^1\Delta$, $^3\Sigma^+$ and $^3\Pi$ symmetries all cross the CO^+ ground state curve near $R = 2.3a_0$. This means that any theoretical study of low-energy dissociative recombination of CO^+ will need to start by considering all these four curves even at the lowest energies.

As mentioned above, states with Σ^- spatial symmetry cannot couple to the target states of the Σ^+ spatial symmetry. This means that the first series of resonance for both $^1\Sigma^-$ and $^3\Sigma^-$ symmetries actually have zero width and are therefore effectively bound states in the continuum. Table 3 gives energies and effective quantum numbers for these states. The data

Table 2. Resonance positions and widths (in Ryd), and effective quantum numbers (ν) at $R = 2.132a_0$ for triplet total symmetries of the CO molecule below the first two CO^+ excited states. Numbers within brackets indicate power of 10.

Below $^2\Pi$ state			Below $^4\Sigma^+$ state		
Position	Width	ν	Position	Width	ν
$^3\Sigma^+$					
0.09116	0.3450 (-03)	2.2540	0.3053	0.5747 (-02)	2.3304
0.1790	0.1572 (-03)	3.0288	0.3143	0.1194 (-02)	2.3895
0.1950	0.1946 (-03)	3.2787	0.3574	0.2796 (-02)	2.7520
0.2216	0.7724 (-03)	3.8817	0.3689	0.1177 (-02)	2.8800
0.2297	0.2507 (-03)	4.1411	0.3822	0.9634 (-03)	3.0541
0.2347	0.1450 (-03)	4.3323	0.4040	0.3654 (-03)	3.4207
0.2525	0.3858 (-04)	5.3090	0.4155	0.3577 (-03)	3.6773
0.2628	0.2102 (-04)	6.3058	0.4226	0.4015 (-03)	3.8664
0.2675	0.3713 (-05)	6.9934	0.4268	0.4016 (-04)	3.9959
			0.4290	0.3768 (-03)	4.0656
			0.4386	0.1709 (-03)	4.4329
			0.4471	0.2020 (-03)	4.8608
			0.4506	0.2287 (-03)	5.0742
			0.4556	0.9297 (-04)	5.4380
$^3\Pi$					
0.0191	0.6342 (-03)	1.9286	0.3025	0.1845 (-02)	2.3129
0.0309	0.4981 (-03)	1.9723	0.3495	0.1257 (-02)	2.6734
0.1133	0.7167 (-03)	2.3925	0.3838	0.6216 (-03)	3.0758
0.1660	0.4748 (-03)	2.8627	0.3994	0.5882 (-03)	3.3326
0.1824	0.3661 (-03)	3.0771	0.4127	0.2164 (-02)	3.6097
0.1859	0.1193 (-02)	3.1303	0.4269	0.6898 (-04)	3.9974
0.1992	0.1251 (-03)	3.3561	0.4297	0.1301 (-03)	4.0910
0.2031	0.2435 (-03)	3.4332	0.4364	0.2987 (-03)	4.3436
0.2205	0.1118 (-03)	3.8493	0.4510	0.9525 (-04)	5.0974
0.2255	0.1138 (-04)	4.0011	0.4544	0.1466 (-03)	5.3425
0.2290	0.7073 (-04)	4.1158	0.4577	0.2995 (-02)	5.6114
0.2371	0.8348 (-04)	4.4330	0.4626	0.4277 (-04)	6.1004
$^3\Delta$					
0.09416	0.4029 (-03)	2.2714	0.3613	0.2080 (-03)	2.7935
			0.4120	0.1415 (-03)	3.5942
			0.4258	0.2046 (-04)	3.9644
			0.4344	0.2562 (-03)	4.2620
			0.4487	0.1175 (-04)	4.9561

in table 3 have been organized to emphasize the near-degeneracy of singlet and triplet states. The small splitting between the singlets and triplets is to be expected for high Rydberg states where spin coupling of the outer electron to the core is weak. The two states, one singlet and one triplet, which are not given as pairs appear to be intruders which will therefore have much lower effective quantum numbers.

3.2. Electronic excitation

Figures 3 and 4 present cross sections for electron impact electronic excitation of CO^+ . These cross sections were calculated by summing the contributions of singlet and triplet




Figure 1. Singlet CO resonance curves as a function of internuclear separation with actual calculated points denoted by stars. The symmetry of each state is indicated in the panel. Also shown are potential energy curves for the three lowest CO⁺ target states of doublet spin symmetry. Resonance which cross the CO⁺ X $^2\Sigma^+$ curve are bound.

symmetries up to and including Δ using the frozen nuclei approximation and a CO⁺ bondlength of $2.132a_0$. Given that the nuclei in CO⁺ move slowly compared to the electron impact the fixed nuclei approximation is probably a reasonable starting point. Both figures show




Figure 2. Triplet CO resonance curves as a function of internuclear separation with actual calculated points denoted by stars. The symmetry of each state is indicated in the panel. Also shown are potential energy curves for the three lowest CO^+ target states. Resonance which cross the $\text{CO}^+ X^2\Sigma^+$ curve are bound.

considerable structure. This structure is associated with the many resonances found in the system. These resonances are, at least at the lower energies considered, real but in practise their influence is washed out by the vibrational motion of the nuclei. This means that no particular significance should be given to the individual peaks and troughs shown in our cross section figures.

Figure 3 gives our predictions for excitation to the first excited state of CO^+ , the $A^2\Pi$ state. As far as we know, there are no previous data on this process for us to compare with. The one previous electron impact excitation measurement concerns electron impact dissociation.

Figure 4 presents our estimate of the electron impact dissociation cross section as a function of electron energy. It is well known that this process goes via electron impact electronic excitation, particularly to dissociative electronic states. In estimating this cross section we have assumed that all electron impact electronic excitation which occurs above the dissociation threshold of 8.5 eV leads to dissociation. This will clearly lead to an overestimate of the dissociation cross section, particularly close to the dissociation threshold. Conversely our calculations only include electronically excited states of CO^+ up to about 12.5 eV. At electron impact energies above this it is likely that our predictions will be an underestimate




Figure 3. Electron impact excitation cross section for $\text{CO}^+ \text{X } ^2\Sigma^+ \rightarrow \text{A } ^2\Pi$ calculated for a fixed bondlength of $R = 2.132a_0$.

Table 3. Energies, in Ryd, and effective quantum numbers for Σ^- bound states in the continuum at $R = 2.132a_0$.

Singlet states		Triplet states	
Position	ν	Position	ν
0.1089	2.3631	0.1087	2.3620
0.1849	3.1138	0.1845	3.1083
0.2002	3.3757	0.2002	3.3753
0.2256	4.0034	0.2256	4.0034
0.2294	4.1306	0.2292	4.1249
0.2359	4.3820	0.2358	4.3791
		0.2463	4.8955
0.2501	5.1373	0.2500	5.1331
0.2535	5.3852	0.2536	5.3910
0.2602	5.9999	0.2603	6.0054
0.2615	6.1407	0.2614	6.1357
0.2635	6.3870	0.2635	6.3898
0.2684	7.1428	0.2684	7.1377
0.2697	7.3881	0.2697	7.3904
0.2729	8.1441	0.2729	8.1390
0.2738	8.3888	0.2738	8.3909
0.2744	8.5667		
0.2760	9.1450	0.2760	9.1398
0.2766	9.3893	0.2767	9.3912

because of this. Despite these caveats our results are comparable to the experimental studies (Mitchell and Hus 1985, Rosen *et al* 1998) of this process. Given that our results will overestimate the amount of dissociation near the dissociation threshold, our results can be taken to be closer to those of Rosen *et al* (1998) than Mitchell and Hus (1985). To improve on our estimate would require both detailed consideration of nuclear motion (see Trevisan and Tennyson (2001)) and inclusion of many more excited states of CO^+ . Such a study is beyond the scope of this work.




Figure 4. Estimated electron impact dissociation cross section for CO^+ calculated for a fixed bondlength of $R = 2.132a_0$. Also shown are the measurements of Mitchell and Hus (1985) (filled circles) and Rosen *et al* (1998) (open diamonds).

4. Conclusions

We have used the UK *R*-matrix codes to consider electron collisions with the CO^+ molecular ion. As is usual for electron collisions with positively charged ions, there are many series of Feshbach resonances associated with these collisions. The low-lying resonances identified are important for studies of dissociative recombination. Previous experience suggests that for a many-electron target ion it is very hard to get the crossing point between such resonances and the ground state of the associated ion precisely (Schneider *et al* 2000). However, it has been shown (Tennyson 1996b, Rabadán and Tennyson 1997, Schneider *et al* 2000) that such error can be corrected for by shifts in the small, geometry independent, quantum defect. An increase in the quantum defect in our calculation by about 0.05 is to be anticipated. Such an approach has been found to both give excellent results when compared with experiment (Schneider *et al* 2000) and to be useful for predictions (Motapon *et al* 2006).

We present data for electron impact electronic excitation of CO^+ . Here our results suggest that the measurements of the electron impact dissociation cross section by Rosen *et al* (1998) is probably closer to the true result than that of Mitchell and Hus (1985). Comparisons of electron impact electronic excitation cross sections between theory and experiment are difficult not only because of the theoretical approximations inherent in the calculations—here, for example, we have used the fixed nuclei approximation—but also because measured electronic excitation cross sections often include cascade effects (Morgan and Tennyson 1993) whereas calculations normally yield the direct excitation cross section only.

Finally we note that our calculations intrinsically contain information on a variety of other processes involving continuum states of CO . It is our plan to use these wavefunctions to study electron impact rotational excitation of CO^+ as part of a study on the process for a variety of astronomical molecules. This can be done with confidence given the recent validation of our method (Faure *et al* 2006).

Acknowledgments

We thank the Royal Society and the Indian National Science Academy for funding KC's visit to UCL during which this work was performed.

References

- Burke P G and Berrington K A (eds) 1993 *Atomic and Molecular Processes, an R-matrix Approach* (Bristol: Institute of Physics Publishing)
- Buttle P J A 1967 *Phys. Rev.* **160** 719–29
- Chakrabarti K and Tennyson J 2006 *J. Phys. B: At. Mol. Opt. Phys.* **39** 1485–97
- Erickson N R, Snell R L, Loren R B, Mundy L and Plambeck R L 1981 *Astrophys. J.* **245** L83–6
- Faure A, Kokoouline V, Greene C H and Tennyson J 2006 *J. Phys. B: At. Mol. Opt. Phys.* **39** 4261–73
- Florescu-Mitchell A I and Mitchel J B A 2006 *Phys. Rep.* **430** 277–374
- Fox J L and Hac A 1999 *J. Geophys. Res.* **104** 24729–37
- Fuente A and Martin Pintado J 1997 *Astrophys. J.* **477** L107
- Gorfinkiel J D, Morgan L A and Tennyson J 2002 *J. Phys. B: At. Mol. Opt. Phys.* **35** 543–55
- Huber K P and Herzberg G (eds) 1979 *Constants of Diatomic Molecules* (New York: Van Nostrand Reinhold)
- Kirby-Docken K and Liu B 1977 *J. Chem. Phys.* **66** 4309–16
- Lecointre J, Belic D S, Cherkani-Hassani H, Jureta J J and Defrance P 2006 *J. Phys. B: At. Mol. Opt. Phys.* **39** 3275–97
- McLaughlin B M, Ballance C P and Berrington K A 2001 *J. Phys. B: At. Mol. Opt. Phys.* **34** L179–86
- McLean A D 1971 *Conference on Potential Energy Surfaces in Chemistry* ed W A Lester Jr (San Jose, CA: IBM Research Laboratory) p 87
- Mitchell J B A and Hus H 1985 *J. Phys. B: At. Mol. Opt. Phys.* **18** 547–55
- Morgan L A 1984 *Comput. Phys. Commun.* **31** 419
- Morgan L A and Tennyson J 1993 *J. Phys. B: At. Mol. Opt. Phys.* **26** 2429–41
- Motapon O, Fifiirig M, Florescu A, Waffeu-Tamo F O, Crumeyrolle O, Varin-Breant G, Bultel A, Vervisch P, Tennyson J and Schneider I F 2006 *Plasma Phys. Sci. Technol.* **15** 23–32
- Noble C J and Nesbet R K 1984 *Comput. Phys. Commun.* **33** 399
- Okada K and Iwata S 2000 *J. Chem. Phys.* **112** 1804
- Rabadan I and Tennyson J 1996 *J. Phys. B: At. Mol. Opt. Phys.* **29** 3747–61
- Rabadán I and Tennyson J 1997 *J. Phys. B: At. Mol. Opt. Phys.* **30** 1975–88
- Rabadán I and Tennyson J 1998 *J. Phys. B: At. Mol. Opt. Phys.* **31** 4485–7 (erratum)
- Rosen S, Peverall R, Larsson M, Le Padellec A, Semaniak J, Larson A, Stromholm C, van der Zande W J, Danared H and Dunn G H 1998 *Phys. Rev. A* **57** 4462–71
- Sarpal B K, Branchett S E, Tennyson J and Morgan L A 1991 *J. Phys. B: At. Mol. Opt. Phys.* **24** 3685–99
- Schneider I F, Rabadán I, Carata L, Tennyson J, Andersen L H and Suzor-Weiner A 2000 *J. Phys. B: At. Mol. Opt. Phys.* **33** 4849–61
- Storzer H, Stutzki J and Sternberg A 1995 *Astron. Astrophys.* **296** L9–L12
- Tennyson J 1996a *J. Phys. B: At. Mol. Opt. Phys.* **29** 1817–28
- Tennyson J 1996b *J. Phys. B: At. Mol. Opt. Phys.* **29** 6185–201
- Tennyson J, Burke P G and Berrington K A 1987 *Comput. Phys. Commun.* **47** 207–12
- Tennyson J and Noble C J 1985 *J. Phys. B: At. Mol. Phys.* **18** 155–65
- Trevisan C S and Tennyson J 2001 *J. Phys. B: At. Mol. Opt. Phys.* **34** 2935–49