

ExoMol line list – XXXIV. A rovibrational line list for phosphinidene (PH) in its $X^3\Sigma^-$ and $a^1\Delta$ electronic states

Jonathan Langleben,¹ Jonathan Tennyson¹,^{1*} Sergei N. Yurchenko¹ and Peter Bernath²

¹Department of Physics and Astronomy, University College London, London WC1E 6BT, UK

²Department of Chemistry and Biochemistry, Old Dominion University, 4541 Hampton Boulevard, Norfolk, VA 23529, USA

Accepted 2019 July 2. Received 2019 June 30; in original form 2019 April 28

ABSTRACT

A rovibronic line list for the ground ($X^3\Sigma^-$) and first excited ($a^1\Delta$) states of phosphinidene, ^{31}PH , is computed. The line list is designed for studies of exoplanetary and cool stellar atmospheres with temperatures up to 4000 K. A combination of empirical and *ab initio* data is used to produce the line list: potential energy curves (PECs) are fitted using experimental transition frequencies; these transitions are reproduced with a root mean square error of 0.01 cm^{-1} . The nuclear Schrödinger equation is solved using these PECs plus Born–Oppenheimer and spin splitting correction terms. Line intensities and Einstein A coefficients are computed using *ab initio* dipole moment curves for X – X and a – a transitions. The resulting LaTY line list, which contains 65 055 transitions for 2528 rovibronic states up to $24\,500\text{ cm}^{-1}$ and $J = 80$, is used to simulate spectra in emission and absorption for a range of temperatures. The line list is made available in electronic form at the CDS and ExoMol data bases.

Key words: molecular data – opacity – astronomical data bases: miscellaneous – planets and satellites: atmospheres – stars: low-mass.

1 INTRODUCTION

The biochemistry of living organisms is heavily dependent on phosphorus. It is important for the storage of genetic information in the form of DNA and RNA (nucleic acids) and is a key contributor to the structure of the cell membrane. The discovery of phosphorus-bearing species in the astrophysical arena is thus thought to be of great significance (Macia 2005; Cui et al. 2017; Zerkle 2018). There have been numerous astrophysical attempts to detect diatomic phosphorus molecules. To date, only PO (Tenenbaum, Woolf & Ziurys 2007), PN (Ziurys 1987), and CP (Guelin et al. 1990) have been identified in the interstellar medium or circumstellar shells. The discovery of PH has so far eluded astronomers (Hollis et al. 1980; Hjalmarsen, Bergman & Biver 2004). However, models of both the interstellar medium (Thorne et al. 1984; Millar 1991) and (exo-) planetary atmospheres (Visscher, Lodders & Fegley 2006) suggest that there are environments where PH should be present in observable quantities. The purpose of this paper is to provide a comprehensive line list for PH to aid in its possible detection and modelling of its spectrum. This line list supplements those of other phosphorus-bearing molecules, namely PN (Yorke et al. 2014), PH_3 (Sousa-Silva et al. 2015), PO, and PS (Prajapat et al. 2017),

produced as part of the ExoMol project (Tennyson & Yurchenko 2012).

The ground electronic state of PH, known as phosphinidene, is of $^3\Sigma^-$ symmetry. It is a singly bonded species so has a lower dissociation energy, of $D_e(\text{PH}) \approx 3.18(3)\text{ eV}$ (Luo 2007; Rumble 2018) (see detailed discussion below), compared to the multiply bonded phosphorus species that have been detected in space: $D_e(\text{PN}) \approx 6.3\text{ eV}$ (Curry, Herzberg & Herzberg 1933), $D_e(\text{PO}) \approx 5.47\text{ eV}$ (Rao, Reddy & Rao 1981), and $D_e(\text{CP}) \approx 5.41\text{ eV}$ (Shi et al. 2012). The spectrum of PH has been well studied in the laboratory (Pearse & Fowler 1930; Ishaque & Pearse 1939; Rostas, Cossart & Bastien 1974; Davies, Russell & Thrush 1975; Di Stefano et al. 1978; Xuan et al. 1978; Ashfold, Dixon & Stickland 1984; Droege & Engelking 1984; Ohashi, Kawaguchi & Hirota 1984; Gustafsson et al. 1985; Ram & Bernath 1987, 1996; Beutel et al. 1996; Hughes & Brown 1997; Klisch et al. 1998; Di Stefano, Lenzi & Ricci 1999; Fitzpatrick et al. 2003) and its electronic structure has been the subject of a number of computational studies (Bruna et al. 1981; Senekowitsch et al. 1986; Park & Sun 1992; Goto & Saito 1993; Fitzpatrick et al. 2002; Jie-Min et al. 2012; Müller & Woon 2013; Gao & Gao 2014). As discussed below, a number of these works form key inputs to the present study.

2 METHOD

Our general methodology for constructing rotation–vibration line

* E-mail: j.tennyson@ucl.ac.uk




Figure 1. *Ab initio* PECs of the lowest electronic states of PH from Gao & Gao (2014) (solid lines) and the refined $X^3\Sigma^-$ and $a^1\Delta$ PECs from this work (dashed).

Table 1. Potential parameters defining empirical X and a PECs according to equation (1). The units are Å and cm^{-1} .

| Parameters | PEC (X) | PEC(a) |
|------------------|---------------|---------------|
| V_e | 0 | 7569 |
| r_e | 1.422179 | 1.423911 |
| D_e | 25700 | 29431 |
| r_{ref} | 1.422179 | 1.423911 |
| p | 4 | 4 |
| N_L | 1 | 1 |
| N_R | 5 | 4 |
| β_0 | 1.7744781772 | 1.6765776631 |
| β_1 | 0.1557401848 | -0.0443775470 |
| β_2 | 0.0119350618 | 1.0241855379 |
| β_3 | 0.4625698020 | -2.3099593616 |
| β_4 | -0.8284568722 | 1.9917675261 |
| β_5 | 0.9631507337 | |

lists is to use available experimental data to characterize the underlying potential energy curve but to use dipole moments computed *ab initio* (see Tennyson 2012; Tennyson & Yurchenko 2017). We follow this approach here. Since PH has a triplet electronic ground state, it is necessary to supplement this approach with spin coupling terms (Tennyson et al. 2016b). All nuclear motion calculations are performed with our general purpose, variational nuclear motion program for diatomic molecules, DUO (Yurchenko et al. 2016). We give details of this procedure in the remainder of this section.

2.1 Potential energy curve

The $X^3\Sigma^-$ and $a^1\Delta$ potential energy curves (PECs) were represented using the extended Morse oscillator (EMO) function as given

by

$$V(r) = V_e + D_e \left[1 - \exp \left(- \sum_{k=0}^N \beta_k \xi_p^k (r - r_e) \right) \right]^2, \quad (1)$$

where D_e is the dissociation energy, ξ_p is the Šurkus variable given by

$$\xi_p = \left(\frac{r^p - r_{\text{ref}}^p}{r^p + r_{\text{ref}}^p} \right), \quad (2)$$

r_e is the corresponding equilibrium bond length, r_{ref} defines the expansion centre for the ξ_p variable (usually taken at r_e), and the integer value p influences how the function extrapolates beyond the data sensitive region. This form allows for extra flexibility in the degree of the polynomial on the left- or on the right-hand sides of the reference position, which is controlled by the parameters $N = N_L$ and N_R , respectively. The empirical parameters V_e and β_k are derived through refinement to experimental data via a least-squares fit, while the dissociation energies D_e are constrained to their experimental asymptotic energies $A_e = V_e + D_e$ (see discussion below).

A set of *ab initio* multireference configuration interaction with Davidson correction (MRCI+ Q) curves for all low-lying states of PH from Gao & Gao (2014) is given in Fig. 1. We used their $X^3\Sigma^-$ and $a^1\Delta$ PECs as starting approximations for our model. As there was sufficient experimental data available for the ground electronic states of PH (Ram & Bernath 1987, 1996; Goto & Saito 1993; Klisch et al. 1998), the refined EMO PEC of fifth order was essentially determined by empirical refinement rather than from the *ab initio* curve of Gao & Gao (2014). For $a^1\Delta$, only the rotational lines for the (0,0) band, $a-X$ electronic transition have been characterized (Beutel et al. 1996). Therefore only V_e and r_e parameters were refined, while all other parameters defining the corresponding EMO PEC of fourth order were fixed to their *ab initio* values. The final set of potential parameters is listed in Table 1.

Table 2. Expansion parameters defining spin–spin (SS), rotational Born–Oppenheimer breakdown (BOB), and spin–rotation (SR) functional forms according to equations (3) and (4). The bond length is in Å, the SS and SR functions are in cm^{-1} , and the BOB expansions are unitless.

| Equation (4) | SS (X) | BOB (X) | Equation (3) | BOB (a) | SR (X) |
|------------------|--------------|---------------|------------------|--------------|---------------|
| r_{ref} | 1.42218 | 1.42218 | r_{ref} | 1.42 | 1.42 |
| γ_2 | 0.8 | 0.8 | p | 4 | 4 |
| γ_4 | 0 | 0 | C_0 | 0.0047811183 | -0.0797377811 |
| p | 2 | 2 | C_∞ | 0 | 0 |
| C_0 | 1.4728278045 | -0.0001063699 | | | |
| C_1 | 1.0957652053 | 0.0089251295 | | | |
| C_2 | 0.4850903021 | -0.3045117774 | | | |
| C_3 | 0 | 0.3645474486 | | | |
| C_4 | 0 | -0.3901575147 | | | |
| C_∞ | 0 | 0 | | | |




Figure 2. Empirical curves of the $X^3\Sigma^-$ state of PH from in this work: spin–spin (cm^{-1}), spin–rotation (cm^{-1}), and Born–Oppenheimer breakdown (unitless).

Table 3. A comparison of experimentally derived vibrational energy levels (in cm^{-1}) for the $X^3\Sigma^-$ state of ^{31}PH from Ram & Bernath (1996) (Obs.) with those calculated using DUO (Calc.).

| v | Obs. | Calc. | Obs. – Calc. |
|-----|----------------|-------------|--------------|
| 1 | 2276.20901(51) | 2276.2061 | 0.00291 |
| 2 | 4465.02033(74) | 4465.0148 | 0.00553 |
| 3 | 6566.15898(88) | 6566.1561 | 0.00288 |
| 4 | 8578.9443(11) | 8578.9506 | -0.0063 |
| 5 | 10502.1949(13) | 10 502.2006 | -0.0057 |

Different couplings and corrections were modelled using the expansion

$$F(r) = (1 - \xi_p) \sum_{k=0}^N C_k z^k + \xi_p C_\infty, \quad (3)$$

where z is either taken as the Šurkus variable $z = \xi_p$ or using the damped-coordinate z given by

$$z = (r - r_{\text{ref}}) e^{-\gamma_2(r-r_{\text{ref}})^2 - \gamma_4(r-r_{\text{ref}})^4} \quad (4)$$

(see also Prajapat et al. 2017; Yurchenko et al. 2018). Here p (integer), C_k , and C_∞ are adjustable parameters. The expansion centre r_{ref} is typically chosen to be close to the equilibrium value of the ground electronic state. To allow for rotational Born–Oppenheimer breakdown (BOB) effects (Le Roy 2017), which

Table 4. A comparison of experimental submillimetre frequencies (in cm^{-1}) for the $X^3\Sigma^-$ state of ^{31}PH from Goto & Saito (1993) and Klisch et al. (1998) (Obs.) with those calculated using DUO (Calc.).

| J' | \pm' | elf' | N' | J'' | \pm'' | elf'' | N'' | Obs. | Calc. | Obs. – Calc. |
|------|--------|--------|------|-------|---------|---------|-------|---------|---------|--------------|
| 0 | + | e | 1 | 1 | - | e | 0 | 14.1191 | 14.1185 | 0.0006 |
| 1 | + | f | 1 | 1 | - | e | 0 | 18.4634 | 18.4623 | 0.0011 |
| 2 | + | e | 1 | 1 | - | e | 0 | 16.4813 | 16.4811 | 0.0002 |
| 1 | - | e | 2 | 1 | + | f | 1 | 30.9308 | 30.9299 | 0.0009 |
| 1 | - | e | 2 | 0 | + | e | 1 | 35.2757 | 35.2738 | 0.0019 |
| 2 | - | f | 2 | 1 | + | f | 1 | 33.6363 | 33.6336 | 0.0027 |
| 2 | - | f | 2 | 2 | + | e | 1 | 35.6146 | 35.6148 | -0.0002 |
| 3 | - | e | 2 | 2 | + | e | 1 | 33.4474 | 33.4445 | 0.0029 |

Table 5. A comparison of experimental IR frequencies (in cm^{-1}) for the $X^3\Sigma^-$ state of ^{31}PH from Ram & Bernath (1987) (Obs.) with those calculated using DUO (Calc.) for a selection of transitions.

| J' | \pm' | elf' | v' | N' | J'' | \pm'' | elf'' | v'' | N'' | Obs. | Calc. | Obs. – Calc. |
|------|--------|--------|------|------|-------|---------|---------|-------|-------|-----------|-----------|--------------|
| 0 | + | e | 1 | 1 | 1 | – | e | 0 | 2 | 2240.4245 | 2240.4228 | 0.0017 |
| 0 | + | e | 2 | 1 | 1 | – | e | 1 | 2 | 2154.0349 | 2154.0248 | 0.0101 |
| 0 | + | e | 3 | 1 | 1 | – | e | 2 | 2 | 2067.3792 | 2067.3730 | 0.0062 |
| 1 | – | e | 1 | 2 | 0 | + | e | 0 | 1 | 2309.9615 | 2309.9625 | –0.0010 |
| 1 | – | e | 1 | 0 | 2 | + | e | 0 | 1 | 2259.7221 | 2259.7249 | –0.0028 |
| 1 | + | f | 1 | 1 | 2 | – | f | 0 | 1 | 2242.0659 | 2242.0688 | –0.0029 |
| 1 | – | e | 1 | 2 | 2 | + | e | 0 | 3 | 2223.9221 | 2223.9240 | –0.0019 |
| 1 | – | e | 2 | 0 | 2 | + | e | 1 | 1 | 2172.8223 | 2172.8137 | 0.0086 |
| 1 | + | f | 2 | 1 | 2 | – | f | 1 | 1 | 2155.6753 | 2155.6634 | 0.0119 |
| 1 | – | e | 2 | 2 | 2 | + | e | 1 | 3 | 2138.0416 | 2138.0324 | 0.0092 |
| 1 | – | e | 3 | 0 | 2 | + | e | 2 | 1 | 2085.6489 | 2085.6481 | 0.0008 |
| 1 | + | f | 3 | 1 | 2 | – | f | 2 | 1 | 2069.0122 | 2069.0003 | 0.0119 |
| 1 | – | e | 3 | 2 | 2 | + | e | 2 | 3 | 2051.8808 | 2051.8782 | 0.0026 |
| ... | | | | | | | | | | | | |
| 5 | + | f | 3 | 5 | 6 | – | f | 2 | 5 | 1999.0172 | 1999.0052 | 0.0120 |
| 5 | – | e | 3 | 6 | 6 | + | e | 2 | 7 | 1980.2734 | 1980.2631 | 0.0103 |
| 5 | – | e | 4 | 4 | 4 | + | e | 3 | 3 | 2068.7136 | 2068.7384 | –0.0248 |
| 5 | + | f | 4 | 5 | 4 | – | f | 3 | 3 | 2081.4480 | 2081.4679 | –0.0199 |
| 5 | – | e | 4 | 6 | 4 | + | e | 3 | 5 | 2093.6318 | 2093.6574 | –0.0256 |
| 5 | – | e | 4 | 4 | 6 | + | e | 3 | 5 | 1931.4560 | 1931.4863 | –0.0303 |
| 5 | + | f | 4 | 5 | 6 | – | f | 3 | 5 | 1913.6358 | 1913.6587 | –0.0229 |
| 5 | – | e | 4 | 6 | 6 | + | e | 3 | 7 | 1895.3895 | 1895.4050 | –0.0155 |
| 5 | – | e | 5 | 4 | 4 | + | e | 4 | 3 | 1977.0756 | 1977.0671 | 0.0085 |
| 5 | + | f | 5 | 5 | 4 | – | f | 4 | 3 | 1989.2557 | 1989.2403 | 0.0154 |
| 5 | – | e | 5 | 6 | 4 | + | e | 4 | 5 | 2000.8809 | 2000.8782 | 0.0027 |
| 6 | + | e | 1 | 5 | 5 | – | e | 0 | 4 | 2352.4659 | 2352.4577 | 0.0082 |
| 6 | – | f | 1 | 6 | 5 | + | f | 0 | 4 | 2366.1885 | 2366.1869 | 0.0016 |
| ... | | | | | | | | | | | | |
| 10 | + | e | 2 | 11 | 11 | – | e | 1 | 12 | 1962.6171 | 1962.6219 | –0.0048 |
| 10 | + | e | 3 | 9 | 9 | – | e | 2 | 8 | 2219.3963 | 2219.3746 | 0.0217 |
| 10 | – | f | 3 | 10 | 9 | + | f | 2 | 8 | 2229.7211 | 2229.7033 | 0.0178 |
| 10 | + | e | 3 | 11 | 9 | – | e | 2 | 10 | 2239.4441 | 2239.4372 | 0.0069 |
| 10 | + | e | 3 | 9 | 11 | – | e | 2 | 10 | 1922.0451 | 1922.0214 | 0.0237 |
| 10 | – | f | 3 | 10 | 11 | + | f | 2 | 10 | 1901.6330 | 1901.6003 | 0.0327 |
| 10 | + | e | 3 | 11 | 11 | – | e | 2 | 12 | 1880.8366 | 1880.8019 | 0.0347 |
| 10 | + | e | 4 | 9 | 9 | – | e | 3 | 8 | 2126.2936 | 2126.3060 | –0.0124 |
| 10 | – | f | 4 | 10 | 9 | + | f | 3 | 8 | 2136.0604 | 2136.0738 | –0.0134 |
| 10 | + | e | 4 | 11 | 9 | – | e | 3 | 10 | 2145.2285 | 2145.2524 | –0.0239 |
| 10 | + | e | 5 | 9 | 9 | – | e | 4 | 8 | 2031.8406 | 2031.8191 | 0.0215 |
| 10 | – | f | 5 | 10 | 9 | + | f | 4 | 8 | 2041.0076 | 2040.9884 | 0.0192 |
| ... | | | | | | | | | | | | |
| 16 | + | e | 3 | 15 | 15 | – | e | 2 | 14 | 2271.8427 | 2271.8199 | 0.0228 |
| 16 | – | f | 3 | 16 | 15 | + | f | 2 | 14 | 2278.3547 | 2278.3273 | 0.0274 |
| 16 | + | e | 3 | 17 | 15 | – | e | 2 | 16 | 2284.1877 | 2284.1849 | 0.0028 |
| 16 | + | e | 4 | 15 | 15 | – | e | 3 | 14 | 2175.3368 | 2175.3445 | –0.0077 |
| 17 | – | e | 1 | 16 | 16 | + | e | 0 | 15 | 2469.8998 | 2469.8935 | 0.0063 |
| 17 | + | f | 1 | 17 | 16 | – | f | 0 | 15 | 2476.8083 | 2476.8153 | –0.0070 |
| 17 | – | e | 1 | 18 | 16 | + | e | 0 | 17 | 2483.0468 | 2483.0661 | –0.0193 |
| 17 | – | e | 1 | 16 | 18 | + | e | 0 | 17 | 1930.5765 | 1930.5700 | 0.0065 |
| 17 | + | f | 1 | 17 | 18 | – | f | 0 | 17 | 1906.7108 | 1906.7114 | –0.0006 |
| 17 | – | e | 1 | 18 | 18 | + | e | 0 | 19 | 1882.5218 | 1882.5467 | –0.0249 |
| 17 | – | e | 2 | 16 | 16 | + | e | 1 | 15 | 2374.4089 | 2374.4329 | –0.0240 |
| 17 | + | f | 2 | 17 | 16 | – | f | 1 | 15 | 2380.8107 | 2380.8373 | –0.0266 |
| 17 | – | e | 2 | 18 | 16 | + | e | 1 | 17 | 2386.5164 | 2386.5774 | –0.0610 |
| 17 | – | e | 3 | 16 | 16 | + | e | 2 | 15 | 2278.3547 | 2278.3206 | 0.0341 |
| 17 | + | f | 3 | 17 | 16 | – | f | 2 | 15 | 2284.1877 | 2284.1579 | 0.0298 |
| 17 | – | e | 3 | 18 | 16 | + | e | 2 | 17 | 2289.3440 | 2289.3366 | 0.0074 |
| 18 | + | e | 1 | 17 | 17 | – | e | 0 | 16 | 2476.8083 | 2476.7978 | 0.0105 |
| 18 | – | f | 1 | 18 | 17 | + | f | 0 | 16 | 2483.0468 | 2483.0481 | –0.0013 |

Table 6. A comparison of experimental $a^1\Delta - X^3\Sigma^-$ (forbidden) frequencies (in cm^{-1}) from Beutel et al. (1996) (Obs.) with those calculated using DUO (Calc.).

| J' | \pm' | elf' | v' | N' | J'' | \pm'' | elf'' | v'' | N'' | Obs. | Calc. | Obs. – Calc. |
|------|--------|--------|------|------|-------|---------|---------|-------|-------|-----------|-----------|--------------|
| 2 | + | e | 0 | 2 | 3 | – | e | 0 | 4 | 7406.0737 | 7406.0727 | 0.0010 |
| 3 | – | e | 0 | 3 | 4 | + | e | 0 | 5 | 7372.6299 | 7372.6427 | –0.0128 |
| 4 | + | e | 0 | 4 | 5 | – | e | 0 | 6 | 7339.3906 | 7339.3921 | –0.0015 |
| 5 | – | e | 0 | 5 | 6 | + | e | 0 | 7 | 7306.3367 | 7306.3324 | 0.0043 |
| 6 | + | e | 0 | 6 | 7 | – | e | 0 | 8 | 7273.4977 | 7273.4805 | 0.0172 |
| 7 | – | e | 0 | 7 | 8 | + | e | 0 | 9 | 7240.8720 | 7240.8559 | 0.0161 |
| 8 | + | e | 0 | 8 | 9 | – | e | 0 | 10 | 7208.5009 | 7208.4794 | 0.0215 |
| 9 | – | e | 0 | 9 | 10 | + | e | 0 | 11 | 7176.3990 | 7176.3726 | 0.0264 |
| 2 | – | f | 0 | 2 | 3 | + | f | 0 | 2 | 7471.0723 | 7471.0815 | –0.0092 |
| 3 | + | f | 0 | 3 | 4 | – | f | 0 | 3 | 7454.5009 | 7454.5073 | –0.0064 |
| 4 | – | f | 0 | 4 | 5 | + | f | 0 | 4 | 7438.0317 | 7438.0375 | –0.0058 |
| 5 | + | f | 0 | 5 | 6 | – | f | 0 | 5 | 7421.6792 | 7421.6832 | –0.0040 |
| 6 | – | f | 0 | 6 | 7 | + | f | 0 | 6 | 7405.4577 | 7405.4557 | 0.0020 |
| 7 | + | f | 0 | 7 | 8 | – | f | 0 | 7 | 7389.3738 | 7389.3667 | 0.0071 |
| 8 | – | f | 0 | 8 | 9 | + | f | 0 | 8 | 7373.4400 | 7373.4278 | 0.0122 |
| 9 | + | f | 0 | 9 | 10 | – | f | 0 | 9 | 7357.6640 | 7357.6514 | 0.0126 |
| 2 | + | e | 0 | 2 | 2 | – | f | 0 | 1 | 7521.4973 | 7521.5055 | –0.0082 |
| 3 | – | e | 0 | 3 | 3 | + | f | 0 | 2 | 7521.6843 | 7521.6904 | –0.0061 |
| 4 | + | e | 0 | 4 | 4 | – | f | 0 | 3 | 7521.9341 | 7521.9375 | –0.0034 |
| 5 | – | e | 0 | 5 | 5 | + | f | 0 | 4 | 7522.2483 | 7522.2477 | 0.0006 |
| 6 | + | e | 0 | 6 | 6 | – | f | 0 | 5 | 7522.6249 | 7522.6217 | 0.0032 |
| 7 | – | e | 0 | 7 | 7 | + | f | 0 | 6 | 7523.0669 | 7523.0608 | 0.0061 |
| 8 | + | e | 0 | 8 | 8 | – | f | 0 | 7 | 7523.5775 | 7523.5663 | 0.0112 |
| 9 | – | e | 0 | 9 | 9 | + | f | 0 | 8 | 7524.1112 | 7524.1400 | –0.0288 |
| 10 | + | e | 0 | 10 | 10 | – | f | 0 | 9 | 7524.7853 | 7524.7841 | 0.0012 |
| 2 | – | f | 0 | 2 | 1 | + | f | 0 | 0 | 7555.1330 | 7555.1391 | –0.0061 |
| 3 | + | f | 0 | 3 | 2 | – | f | 0 | 1 | 7572.1114 | 7572.1144 | –0.0030 |
| 4 | – | f | 0 | 4 | 3 | + | f | 0 | 2 | 7589.1204 | 7589.1206 | –0.0002 |
| 5 | + | f | 0 | 5 | 4 | – | f | 0 | 3 | 7606.1501 | 7606.1477 | 0.0024 |
| 6 | – | f | 0 | 6 | 5 | + | f | 0 | 4 | 7623.1921 | 7623.1862 | 0.0059 |
| 7 | + | f | 0 | 7 | 6 | – | f | 0 | 5 | 7640.2333 | 7640.2267 | 0.0066 |
| 8 | – | f | 0 | 8 | 7 | + | f | 0 | 6 | 7657.2646 | 7657.2604 | 0.0042 |
| 9 | + | f | 0 | 9 | 8 | – | f | 0 | 7 | 7674.2851 | 7674.2784 | 0.0067 |
| 10 | – | f | 0 | 10 | 9 | + | f | 0 | 8 | 7691.2735 | 7691.2726 | 0.0009 |
| 11 | + | f | 0 | 11 | 10 | – | f | 0 | 9 | 7708.2094 | 7708.2353 | –0.0259 |
| 12 | – | f | 0 | 12 | 11 | + | f | 0 | 10 | 7725.1395 | 7725.1594 | –0.0199 |
| 2 | + | e | 0 | 2 | 1 | – | e | 0 | 0 | 7573.5937 | 7573.6014 | –0.0077 |
| 3 | – | e | 0 | 3 | 2 | + | e | 0 | 1 | 7607.7241 | 7607.7292 | –0.0051 |
| 4 | + | e | 0 | 4 | 3 | – | e | 0 | 2 | 7641.7120 | 7641.7149 | –0.0029 |
| 5 | – | e | 0 | 5 | 4 | + | e | 0 | 3 | 7675.6392 | 7675.6403 | –0.0011 |
| 6 | + | e | 0 | 6 | 5 | – | e | 0 | 4 | 7709.5131 | 7709.5125 | 0.0006 |
| 7 | – | e | 0 | 7 | 6 | + | e | 0 | 5 | 7743.3243 | 7743.3220 | 0.0023 |
| 8 | + | e | 0 | 8 | 7 | – | e | 0 | 6 | 7777.0560 | 7777.0542 | 0.0018 |
| 9 | – | e | 0 | 9 | 8 | + | e | 0 | 7 | 7810.6816 | 7810.6924 | –0.0108 |
| 10 | + | e | 0 | 10 | 9 | – | e | 0 | 8 | 7844.1942 | 7844.2194 | –0.0252 |
| 10 | – | f | 0 | 10 | 11 | + | f | 0 | 10 | 7342.0698 | 7342.0500 | 0.0198 |

become important for high J , the vibrational kinetic energy operator

for each state was extended by

$$-\frac{\hbar^2}{2\mu r^2} \rightarrow -\frac{\hbar^2}{2\mu r^2} (1 + g^{\text{BOB}}(r)), \quad (5)$$

where the unitless X and a BOB functions were represented by the (unitless) polynomial $g^{\text{BOB}}(r) = F(r)$. The X -state BOB function required a fourth-order expansion in terms of damped- z in equation (4), while for the a -state only one C_0 constant from equation (3) was needed. When fitting to the experimental frequencies, we had to include other X -state correction terms, such as spin–spin and spin–rotation (see Yurchenko et al. 2016), for

which the same z -damped expression equation (3) was used, of third and zeroth orders, respectively. All expansion parameters are given in Table 2 and in the supplementary material as part of the DUO input file.

Experimental data were taken primarily from four sources. To determine the X -state curves, submillimetre-wave measurements of the ($N = 0, 1$) rotational spectrum by Goto & Saito (1993) and the ($N = 1, 2$) spectrum by Klisch et al. (1998) were combined with frequencies from an infrared (IR) Fourier transform spectrometer vibration–rotation spectrum of the ground state of PH by Ram & Bernath (1987). The infrared study observed five vibrational bands (1–0, 2–1, 3–2, 4–3, and 5–4) up to a maximum rotation state of $J = 21$. The X -state data set comprised 381 lines split among six fine-structure resolved branches: R_1, R_2, R_3, P_1, P_2 , and P_3 that characterize the triplet pattern arising from the splitting of the lines from the electronic spin angular momentum along the internuclear axis (i.e. different spin projections $\Sigma = -1, 0, 1$; note that the X state is Hund’s case (b) so these projections on the internuclear axis are not good quantum numbers). These data allowed the lowest six vibrational states ($v = 0 \dots 5$) to be characterized. The hyperfine structure of the submillimetre-wave frequencies (Goto & Saito 1993; Klisch et al. 1998) was averaged. For the a -state, 64 a - X IR transition frequencies (0–0 band) from Beutel et al. (1996) were used.

According to Rumble (2018) (with the original reference to Luo 2007), the dissociation value for PH at $T = 298$ K is 297.0 ± 2.1 kJ mole $^{-1}$ or 293.2 kJ mole $^{-1}$ at 0 K (24516 ± 175 cm $^{-1}$ or $3.04(2)$ eV). Using a zero-point energy value of 1170 cm $^{-1}$ from our calculations ($ZPE = D_e - D_0$), this corresponds to $D_e = 25\,687 \pm 175$ cm $^{-1}$ (3.19 eV). The value $A_e(X) = D_e(X) = 25\,700$ cm $^{-1}$ was adopted as the dissociation energy of the X -state in our calculations. Throughout the refinement phase, the equilibrium distance r_e was kept fixed at $r_e = 1.4221$ Å, as spectroscopically determined by Ram & Bernath (1987).

The dissociation channel for the a states can be estimated using the phosphorus atom excitation energy ($^2D_{3/2}$): $A_e(a) = D_e(X) + 11361.02 \approx 37000$ cm $^{-1}$ (Kramida et al. 2019), which was used to constrain the dissociation energy of $a^1\Delta$ while varying the corresponding value of the origin V_e .

The PECs and other empirical curves fitted directly to experimental-measured frequencies ($J \leq 21$) achieved a root-mean-square error of 0.01 cm $^{-1}$. The final refined PECs are shown in Fig. 1 and empirical curves are shown in Fig. 2. Some of the residuals are illustrated in Tables 3–6.

2.2 *Ab initio* dipole moment curves

PH has a permanent dipole and can be represented by a dipole moment curve (DMC) that shows the variation of the dipole with internuclear separation. Gao & Gao (2014) computed an *ab initio* DMC for the $X^3\Sigma^-$ state of PH using an aug-ccpV5Z basis set at the MRCI+ Q level of theory. We used the same method to compute the DMC for the $a^1\Delta$ state with the MOLPRO program (Werner et al. 2012). These DMCs are illustrated in Fig. 3. As expected for a neutral species, the dipole moments tend to zero at large bond lengths.

The $X^3\Sigma^-$ state *ab initio* dipole moment has an equilibrium value of 0.4771 D. This compares reasonably with other theoretical study equilibrium values. Müller & Woon (2013) calculated dipole moments at a number of theoretical levels, using a variety of basis sets. An AV5Z basis set at the coupled cluster (CCSD(T)) level of

theory produced a dipole moment of 0.4410 D. Other studies at the configuration–interaction (CI) level calculated the dipole moment as 0.431 D (Senekowitsch et al. 1986) and 0.432 D (Park & Sun 1992). Meyer & Rosmus (1975) using the coupled electron pair approach (CEPA) produced a dipole moment of 0.481 D, which is closest to the value of Gao & Gao (2014) used for our line list calculations.

The DMCs were represented analytically using the damped- z expansion given by equation (3). This was done in order to reduce the numerical noise in the calculated intensities for high overtones; see recommendations by Medvedev et al. (2016). These DMCs and the empirically defined curves constitute our spectroscopic model, which was then used with DUO to produce a PH line list. The corresponding expansion parameters are listed in Table 7. They can also be found in the supplementary data in the form of the DUO input files together with the corresponding grid representations.

3 RESULTS

3.1 Line list

The spectroscopic model described in the previous sections was used to generate the line list for the ground (X) and first excited (a) electronic states of ^31PH . The LaTY line list was computed using the empirical PECs and correction curves and *ab initio* DMCs described above. All vibrational states up to $v = 17$ (X) and $v = 9$ (a), rotational states up to $J = 80$ and energies up to $D_0(X)$ were considered. A dipole moment cut-off of 10^{-7} D was applied. The zero-point energy was calculated to be 1170.47 cm $^{-1}$ defined as the lowest state of the system ($J = 1, v = 0, N = 0, e(-)$) above the minimum of the X potential ($V_e = 0$ cm $^{-1}$). The final line list contains 2528 states and 65 055 transitions. These levels are sufficient to represent the ground state up to temperatures of about 4000 K, but at these temperatures electronically excited states should also be occupied giving rise to further transitions not considered here. The line list consists of the electric dipole transitions only. Moreover, the $X^3\Sigma^-$ and $a^1\Delta$ states are fully uncoupled and therefore the line list does not include the weak, forbidden $a - X$ transitions observed in Beutel et al. (1996).

The results are provided in the standard ExoMol format (Tennyson et al. 2016a) as states and transitions files, see extracts given in Tables 8 and 9, respectively. Since DUO works in Hund’s case (a) but PH is a Hund’s case (b) molecule for the X state, the ($\pm, \Lambda, \Sigma, \Omega$) quantum numbers were converted to N, elf . For the $a^1\Delta$ state, $N = J, \Sigma = 0, \Lambda = \Omega = 2$. Here N is the rotational quantum number defined as a projection of $N = J - S$, where J and S are the total and spin angular momenta, respectively. The states file also gives Landé g -factors for the various states (Semenov, Yurchenko & Tennyson 2017) that can be used to model the behaviour of these states in a weak magnetic field.

3.2 Partition function

The partition function, $Q(T)$, was calculated by summing the energy levels given by DUO for temperatures up to $T = 4000$ K. ExoMol follows the HITRAN convention (Gamache et al. 2017) of explicitly including the full atomic nuclear spin in the molecular partition function via the nuclear spin statistical number g_{ns} . Since both P and H have nuclear spins $1/2$, $g_{\text{ns}} = 4$.

The partition function at a range of temperatures is catalogued in Table 10. It was compared with sources where the partition function $Q(T)$ was deduced from polynomial approximations (Irwin 1981;




Figure 3. *Ab initio* PH MRCI+*Q*/AV5Z dipole moment curve (DMC) used in this work: the $X^3\Sigma^-$ state DMC is from Gao & Gao (2014), and $a^1\Delta$ DMC was computed as part of this study.

Table 7. Expansion parameters defining X and a DMCs according to equation (4). The units are Å (Angstrom) and D (Debye).

| Parameters | DMC(X) | DMC(a) |
|------------------|---------------|---------------|
| r_{ref} | 1.4222 | 1.4222 |
| γ_2 | 0.1695549437 | 0.1233507271 |
| γ_4 | 0.0116033136 | 0.0210533244 |
| p | 6 | 6 |
| D_0 | 0.4766434841 | 0.4747489601 |
| D_1 | -0.2603505526 | -0.3886599811 |
| D_2 | -1.6119565470 | -1.3959503273 |
| D_3 | -4.4208750252 | -3.5580088368 |
| D_4 | -3.6843015061 | -2.7131260560 |
| D_∞ | 0 | 0 |

Sauval & Tatum 1984; Barklem & Collet 2016). The cited partition functions were all multiplied by 4 to match with the HITRAN convention adopted. It can be seen that all the sources approximately agree with each other between 1000 and 4000 K. Below 1000 K, polynomial representations of $Q(T)$ used by Sauval & Tatum (1984) and Irwin (1981) are not valid; our results are much closer to the modern values of Barklem & Collet (2016) albeit slightly higher probably because of our full treatment of electron spin effects. Above 4000 K our values for $Q(T)$ are lower than those of Sauval & Tatum (1984) and Barklem & Collet (2016); these works include the contribution from electronically excited states that we neglect.

3.3 Experimental spectra

Two vibration–rotation emission spectra were recorded with the Fourier transform spectrometer at the National Solar Observatory

at Kitt Peak in Arizona. The first spectrum (Spectrum 1 shown as the upper red trace in Fig. 4; Ram & Bernath 1987) was recorded with an electrodeless quartz discharge tube excited with a 2450-MHz microwave oscillator. A mixture of 0.45 Torr of hydrogen and 0.04 Torr of white phosphorus vapour flowed through the cell. The spectral resolution was 0.02 cm^{-1} and covered the $1800\text{--}9000 \text{ cm}^{-1}$ region. The second spectrum (Spectrum 2 shown as the lower green trace in Fig. 4) differed only in the gas mixture used: 2.75 Torr of helium, 0.04 Torr of white phosphorus, and 0.03 Torr of methane. Both spectra contained many molecules including CO, CH, PH, CP, P_2 , ArH, CN, and C_2 . The strongest interfering molecule overlapping PH is CO with its $\Delta v = 1$ emission lines. The CO lines were used for wavenumber calibration. The vibration–rotation emission lines of PH are stronger in the first spectrum but the number of interfering lines is somewhat reduced in the second spectrum. Norton–Beer strong apodization was used because the lines were not resolved at 0.02 cm^{-1} resolution and the lines still have residual ‘ringing’ from the instrument line shape function.

3.4 Theoretical spectra

Fig. 5 gives an overview of full PH spectrum in absorption at two temperatures, 300 and 2000 K. At the lower temperature a clear progression of vibrational bands can be seen that is substantially washed out at the higher temperature due to the presence of many more weak lines in the spectrum.

Fig. 6 shows the microwave spectrum of the ground state of PH at $T = 296 \text{ K}$. It is compared with the Cologne Database for Molecular Spectroscopy (CDMS) spectrum (Müller et al. 2005) ($v = 0$), which we have averaged over the hyperfine components. The dipole moment used by CDMS is 0.396 D at r_e (Müller, unpublished work), whereas the dipole moment from this work is 0.477 D at equilibrium

Table 8. Sample extract of the states file for ^{31}PH .

| i | \bar{E} | g_i | J | g | \pm | elf | State | v | N |
|-----|--------------|-------|-----|----------|-------|-------|----------|-----|-----|
| 73 | 16.481110 | 20 | 2 | 1.00004 | + | e | X3Sigma- | 0 | 1 |
| 74 | 100.157012 | 20 | 2 | -0.66632 | + | e | X3Sigma- | 0 | 3 |
| 75 | 2292.186846 | 20 | 2 | 0.99997 | + | e | X3Sigma- | 1 | 1 |
| 76 | 2373.329250 | 20 | 2 | -0.66625 | + | e | X3Sigma- | 1 | 3 |
| 77 | 4480.481941 | 20 | 2 | 0.99989 | + | e | X3Sigma- | 2 | 1 |
| 78 | 4559.093532 | 20 | 2 | -0.66618 | + | e | X3Sigma- | 2 | 3 |
| 79 | 6581.109731 | 20 | 2 | 0.99981 | + | e | X3Sigma- | 3 | 1 |
| 80 | 6657.178701 | 20 | 2 | -0.66610 | + | e | X3Sigma- | 3 | 3 |
| 81 | 7573.601437 | 20 | 2 | 0.66667 | + | e | a1Delta | 0 | 2 |
| 82 | 8593.416817 | 20 | 2 | 0.99974 | + | e | X3Sigma- | 4 | 1 |
| 83 | 8666.923664 | 20 | 2 | -0.66602 | + | e | X3Sigma- | 4 | 3 |
| 84 | 9899.143295 | 20 | 2 | 0.66667 | + | e | a1Delta | 1 | 2 |
| 85 | 10516.156390 | 20 | 2 | 0.99966 | + | e | X3Sigma- | 5 | 1 |
| 86 | 10587.067513 | 20 | 2 | -0.66594 | + | e | X3Sigma- | 5 | 3 |

Note. i : state counting number.

\bar{E} : state energy in cm^{-1} .

g_i : total statistical weight, equal to $g_{\text{ns}}(2J + 1)$.

J : total angular momentum.

g : Landé g -factors.

elf : rotationless parity.

\pm : total parity.

State: electronic state.

v : state vibrational quantum number.

N : rotational quantum number; $N = J - S$ (S is spin).

Table 9. Sample extract of the transitions file for ^{31}PH .

| f | i | A_{fi} | $\bar{\nu}_{fi}$ |
|------|------|------------|------------------|
| 100 | 134 | 2.1325E-06 | 31.942920 |
| 162 | 135 | 3.0255E-05 | 31.959112 |
| 102 | 60 | 2.8977E-05 | 32.215418 |
| 256 | 285 | 3.2537E-06 | 32.223799 |
| 44 | 4 | 1.7832E-04 | 32.237083 |
| 174 | 75 | 6.7300E-04 | 32.439634 |
| 121 | 79 | 8.4497E-05 | 32.564436 |
| 119 | 20 | 5.1188E-04 | 32.625072 |
| 2047 | 2069 | 1.1922E-11 | 32.678476 |

Note. f : state number of final state.

i : state number of initial state.

A_{fi} : Einstein A coefficient.

$\bar{\nu}_{fi}$: transition wavenumber.

or, more importantly, our $v = 0$ state vibrationally averaged dipole is 0.4499 D. This disparity can be seen in the graph, with our lines being more intense due to the larger dipole moment. Partition function values from CDMS agree with that from this work. At 300 K, the CDMS value was 302.12, while our value is 302.14. The line strength depends on the dipole moment squared. With this in mind, our intensities are about 1.27 times higher than those given by CDMS. We suggest that CDMS may wish to rescale their intensity values to ours. Conversely, the CDMS frequencies rely directly on data taken in the same frequency region and include the hyperfine components and must be considered more accurate than ours in this region; they should be used for any attempts to detect PH in the interstellar medium.

Fig. 4 compares the IR rovibrational experimental spectra from Ram & Bernath (1987) with calculations from this work. We

assumed the non-local thermodynamic equilibrium (LTE) model based on two temperatures, rotational T_{rot} and vibrational T_{vib} , as implemented in DUO. The temperatures were adjusted to match the experimental emission spectra of PH, with the final values of 800 and 2300 K, respectively. The grey vertical line indicates the experimental line positions from Ram & Bernath (1987). The good agreement with the observed spectrum of Ram & Bernath (1987) is found across the entire region and confirms the accuracy of our calculations.

4 CONCLUSION

A comprehensive line list for the ground (X) and first excited (a) electronic states of ^{31}PH , known as LaTY, is presented. It is based on an accurate PECs, BOB, SS, and SR curves obtained by fitting to a set of experimental transition line frequencies and extrapolating to higher rovibrational levels and *ab initio* DMCS. Future work can include analysis and investigations of higher electronic states of PH with the aim of creating a line list appropriate at shorter wavelengths. Previous studies on the low-lying $b^1\Sigma^+$ of PH (Di Stefano et al. 1978, 1999; Xuan et al. 1978) and on the strongly dipole-allowed

Table 10. Comparison with calculated partition functions, $Q(T)$, of Sauval & Tatum (1984), Irwin (1981), and Barklem & Collet (2016).

| T (K) | This work | Sauval & Tatum | Irwin | Barklem & Collet |
|---------|-----------|----------------|---------|------------------|
| 100 | 103.24 | 112.20 | 12.03 | 103.33 |
| 200 | 202.58 | 222.11 | 125.10 | 202.68 |
| 300 | 302.14 | 320.89 | 261.37 | 302.25 |
| 400 | 401.99 | 416.37 | 383.16 | |
| 500 | 502.54 | 512.03 | 494.35 | 502.64 |
| 600 | 604.56 | 609.72 | 600.96 | |
| 700 | 708.99 | 710.52 | 706.92 | 709.08 |
| 800 | 816.74 | 815.16 | 814.60 | |
| 900 | 928.57 | 924.17 | 925.49 | |
| 1000 | 1045.12 | 1037.97 | 1040.52 | 1045.20 |
| 1500 | 1713.31 | 1689.58 | 1694.33 | 1713.48 |
| 2000 | 2549.16 | 2501.19 | 2499.05 | 2550.26 |
| 2500 | 3580.45 | 3500.15 | 3467.86 | |
| 3000 | 4834.19 | 4713.74 | 4609.02 | 4847.12 |
| 4000 | 8112.16 | 7901.37 | 7436.84 | 8180.60 |

A $^3\Pi - X^3\Sigma^-$ system (Fitzpatrick et al. 2002, 2003; Gao & Gao 2014) in the near-ultraviolet provide a good starting point for future analysis of the PH molecule.

The line list can be downloaded from the CDS, via <ftp://cdsarc.u-strasbg.fr/pub/cats/J/MNRAS/>, or <http://cdsarc.u-strasbg.fr/viz-bin/qcat?J/MNRAS/>, or from www.exomol.com.

ACKNOWLEDGEMENTS

This work was supported by the UK Science and Technology Research Council (STFC) No. ST/R000476/1 and the COST action MOLIM No. CM1405. This work made extensive use of UCL's Legion high performance computing facility along with the STFC DiRAC HPC facility supported by BIS National E-infrastructure capital grant ST/J005673/1 and STFC grants ST/H008586/1 and ST/K00333X/1. Some support was provided by the NASA Laboratory Astrophysics Program.

REFERENCES

Ashfold M. N. R., Dixon R. N., Stickland R. J., 1984, *Chem. Phys. Lett.*, 111, 226
 Barklem P. S., Collet R., 2016, *A&A*, 588, A96
 Beutel M., Setzer K. D., Shestakov O., Fink E. H., 1996, *Chem. Phys. Lett.*, 249, 183
 Bruna P. J., Hirsch G., Peyerimhoff S. D., Buenker P. J., 1981, *Mol. Phys.*, 42, 875
 Cui D., Tian F., Wang Y., Li C., Yu C., Yu L., 2017, *Astrobiology*, 17, 1219
 Curry J., Herzberg L., Herzberg G., 1933, *J. Chem. Phys.*, 1, 749
 Davies P. B., Russell D. K., Thrush B. A., 1975, *Chem. Phys. Lett.*, 36, 280
 Di Stefano G., Lenzi M., Margani A., Xuan C. N., 1978, *J. Chem. Phys.*, 68, 959
 Di Stefano G., Lenzi M., Ricci A., 1999, *Chem. Phys. Lett.*, 246, 267
 Droge A., Engelking P., 1984, *J. Chem. Phys.*, 80, 5926
 Fitzpatrick J. A. J., Chekhlov O. V., Morgan D. R., Burrows R. W., Western C. M., 2002, *Phys. Chem. Chem. Phys.*, 4, 1114
 Fitzpatrick J. A. J., Chekhlov O. V., Western C. M., Ashworth S. H., 2003, *J. Chem. Phys.*, 118, 4539
 Gamache R. R. et al., 2017, *J. Quant. Spectrosc. Radiat. Transf.*, 203, 70

Gao Y., Gao T., 2014, *Spectrochim. Acta A*, 118, 308
 Goto M., Saito S., 1993, *Chem. Phys. Lett.*, 211, 443
 Guelin M., Cernicharo J., Paubert G., Turner B. E., 1990, *A&A*, 230, L9
 Gustafsson O., Kindvall G., Larsson M., Senekowitsch J., Sigraý P., 1985, *Mol. Phys.*, 56, 1369
 Hjalmarson A., Bergman P., Biver N., 2004, *Adv. Space Res. J.*, 36, 1031
 Hollis J. M., Snyder L. E., Lovas F. J., Ulich B. L., 1980, *ApJ*, 241, 158
 Hughes R. A., Brown J. M., 1997, *J. Mol. Spectrosc.*, 185, 197
 Irwin A. W., 1981, *ApJS*, 45, 621
 Ishaque M., Pearse R. W. B., 1939, *Proc. R. Soc. Lond. Ser. A*, 173, 0265
 Jie-Min W., Jin-Feng S., De-Heng S., Zun-Lue Z., Wen-Tao L., 2012, *Acta Phys. Sin.*, 61, 063104
 Klisch E., Klein H., Winnewisser G., Herbst E., 1998, *Z. Naturforschung A*, 53, 733
 Kramida A., Ralchenko Yu., Reader J., NIST ASD Team, 2019, NIST Atomic Spectra Database (ver. 5.6.1). National Institute of Standards and Technology, Gaithersburg, MD [Online]. Available: <https://physics.nist.gov/asd> [2019 April 13]
 Le Roy R. J., 2017, *J. Quant. Spectrosc. Radiat. Transf.*, 186, 167
 Luo Y. R., 2007, *Comprehensive Handbook of Chemical Bond Energies*, 1st edn. CRC Press, Boca Raton, FL
 Macia E., 2005, *Chem. Soc. Rev.*, 34, 691
 Medvedev E. S., Meshkov V. V., Stolyarov A. V., Ushakov V. G., Gordon I. E., 2016, *J. Mol. Spectrosc.*, 330, 36
 Meyer W., Rosmus P., 1975, *J. Chem. Phys.*, 63, 2356
 Millar T. J., 1991, *A&A*, 242, 241
 Müller H. S. P., Woon D. E., 2013, *J. Phys. Chem. A*, 117, 13868
 Müller H. S. P., Schlöder F., Stutzki J., Winnewisser G., 2005, *J. Mol. Structure*, 742, 215
 Ohashi N., Kawaguchi K., Hirota E., 1984, *J. Mol. Spectrosc.*, 103, 337
 Park J. K., Sun H., 1992, *Chem. Phys. Lett.*, 195, 469
 Pearse R. W. B., Fowler A., 1930, *Proc. R. Soc. Lond. Ser. A*, 129, 328
 Prajapat L., Jagoda P., Lodi L., Gorman M. N., Yurchenko S. N., Tennyson J., 2017, *MNRAS*, 472, 3648
 Ram R. S., Bernath P. F., 1987, *J. Mol. Spectrosc.*, 122, 275
 Ram R. S., Bernath P. F., 1996, *J. Mol. Spectrosc.*, 176, 329
 Rao T. V. R., Reddy R. R., Rao P. S., 1981, *Physica B+C*, 106, 445
 Rostas J., Cossart D., Bastien J. R., 1974, *Can. J. Phys.*, 52, 1274
 Rumble J., 2018, *CRC Handbook of Chemistry and Physics*, 99th edn. CRC Press, Boca Raton, FL
 Sauval A. J., Tatum J. B., 1984, *ApJS*, 56, 193




Figure 4. Experimental infrared spectra (see text for details) of the $X^3\Sigma^-$ state of PH compared to the ExoMol emission (photons s^{-1}) at $T_{\text{rot}} = 800$ K and $T_{\text{vib}} = 2300$ K using a Voigt line profile with $\gamma = 0.15$ cm^{-1} . Vertical grey lines indicate experimental line positions from Ram & Bernath (1987).

Semenov M., Yurchenko S. N., Tennyson J., 2017, *J. Mol. Spectrosc.*, 330, 57
 Senekowitsch J., Rosmus P., Werner H. J., Larsson M., 1986, *Z. Naturforschung A*, 41, 719
 Shi D., Xing W., Sun J., Zhu Z., Liu Y., 2012, *J. Mol. Spectrosc.*, 276, 1
 Sousa-Silva C., Al-Refaie A. F., Tennyson J., Yurchenko S. N., 2015, *MNRAS*, 446, 2337
 Tenenbaum E. D., Woolf N. J., Ziurys L. M., 2007, *ApJ*, 666, L29
 Tennyson J., 2012, *WIREs Comput. Mol. Sci.*, 2, 698
 Tennyson J., Yurchenko S. N., 2012, *MNRAS*, 425, 21
 Tennyson J., Yurchenko S. N., 2017, *Mol. Astrophys.*, 8, 1
 Tennyson J. et al., 2016a, *J. Mol. Spectrosc.*, 327, 73
 Tennyson J., Lodi L., McKemmish L. K., Yurchenko S. N., 2016b, *J. Phys. B: At. Mol. Opt. Phys.*, 49, 102001

Thorne L. R., Anicich V. G., Prasad S. S., Huntress W. T., Jr, 1984, *ApJ*, 280, 139
 Visscher C., Lodders K., Fegley B., Jr, 2006, *ApJ*, 648, 1181
 Werner H.-J., Knowles P. J., Knizia G., Manby F. R., Schütz M., 2012, *WIREs Comput. Mol. Sci.*, 2, 242
 Xuan C. N., di Stefano G., Lenzi M., Margani A., Mele A., 1978, *Chem. Phys. Lett.*, 57, 207
 Yorke L., Yurchenko S. N., Lodi L., Tennyson J., 2014, *MNRAS*, 445, 1383
 Yurchenko S. N., Lodi L., Tennyson J., Stolyarov A. V., 2016, *Comput. Phys. Commun.*, 202, 262
 Yurchenko S. N., Sinden F., Lodi L., Hill C., Gorman M. N., Tennyson J., 2018, *MNRAS*, 473, 5324
 Zerkle A. L., 2018, *Philos. Trans. R. Soc. A*, 376, 20170401




Figure 5. Absorption spectrum of PH at different temperatures computed using ExoMol line list and a Gaussian line profile of 2 cm^{-1} . The contribution from the a -band at $T = 2000 \text{ K}$ is shown separately.




Figure 6. Pure rotational band within $v = 0$ for the rovibrational absorption stick spectra of the $X^3\Sigma^-$ state of PH at 296 K.

Ziurys L. M., 1987, *ApJ*, 321, L81

SUPPORTING INFORMATION

Supplementary data are available at [MNRAS](https://www.mnras.org/) online.

PH.pf
PH.states

PH_model.inp

Please note: Oxford University Press is not responsible for the content or functionality of any supporting materials supplied by the authors. Any queries (other than missing material) should be directed to the corresponding author for the article.

This paper has been typeset from a $\text{\TeX}/\text{\LaTeX}$ file prepared by the author.