

RESEARCH ARTICLE OPEN ACCESS

MARVEL Analysis of High-Resolution Rovibrational Spectra of $^{16}\text{O}^{13}\text{C}^{18}\text{O}$ Ala'a A. A. Azzam¹ | Jonathan Tennyson² | Sergei N. Yurchenko² | Tibor Furtenbacher³ | Attila G. Császár^{3,4}¹Department of Physics, The University of Jordan, Amman, Jordan | ²Department of Physics and Astronomy, University College London, London, UK | ³HUN-REN-ELTE Complex Chemical Systems Research Group, Budapest, Hungary | ⁴Institute of Chemistry, ELTE Eötvös Loránd University, Budapest, Hungary**Correspondence:** Ala'a A. A. Azzam (alaa.azzam@ju.edu.jo) | Jonathan Tennyson (j.tennyson@ucl.ac.uk)**Received:** 30 September 2024 | **Accepted:** 18 October 2024**Funding:** The authors thank STFC for funding the UK – Jordan collaboration under the Newton Fund grant ST/T001429/1. JT acknowledges the support of the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation program through Advance Grant number 883830. The work in Budapest has received funding from the HUN-REN Hungarian Research Network and the National Research, Development and Innovation Office (NKFIH, grant no. K138233). This publication supports research performed within the COST Action CA21101 "Confined molecular systems: from a new generation of materials to the stars" (COSY), funded by the European Cooperation in Science and Technology (COST). Ala'a Azzam thanks AstroJo Institute for providing the necessary support to complete this research project.**Keywords:** $^{16}\text{O}^{13}\text{C}^{18}\text{O}$ | carbon dioxide | high-resolution spectroscopy | line positions | MARVEL analysis | rovibrational energy levels

ABSTRACT

A large set of validated experimental transitions and empirical rovibrational energy levels are reported for the fifth most abundant carbon dioxide isotopologue, $^{16}\text{O}^{13}\text{C}^{18}\text{O}$ (in a shorthand notation, 638). Validation of the transitions and determination of the empirical energy levels are based on a compiled and carefully checked dataset, collected from 35 literature sources, containing 12 348/7432 measured/unique lines in the wavenumber range of 578–9318 cm^{-1} . The MARVEL (Measured Active Rotational-Vibrational Energy Levels) protocol, built upon the theory of spectroscopic networks, not only validates the vast majority of the measured transitions, but also yields 3975 empirical rovibrational energy levels, with uncertainty estimates compliant with the experimental uncertainties of the transitions.

1 | Introduction

Recently, we decided to construct an extensive database of empirical rovibrational energy levels, based on spectral features measured under high resolution, for all the isotopologues of carbon dioxide involving the ^{12}C , ^{13}C , ^{16}O , ^{17}O , and ^{18}O isotopes. The present contribution extends the list of CO_2 isotopologues treated by us, $^{16}\text{O}^{13}\text{C}^{16}\text{O}$ (636, according to a shorthand notation following HITRAN [1] parlance) [2], $^{16}\text{O}^{12}\text{C}^{18}\text{O}$ (628) [3], $^{18}\text{O}^{12}\text{C}^{18}\text{O}$ (828) [4], $^{18}\text{O}^{13}\text{C}^{18}\text{O}$ (838) [4], and $^{17}\text{O}^{12}\text{C}^{18}\text{O}$ (728) [4]. Investigation of the parent isotopologue, $^{16}\text{O}^{12}\text{C}^{16}\text{O}$ (626), is also under way by our laboratories [5]. With a natural abundance of

0.004434%, the carbon dioxide isotopologue $^{16}\text{O}^{13}\text{C}^{18}\text{O}$ (638) is the fifth most abundant isotopologue of CO_2 [1]. Similar to the other projects, the empirical rovibrational energy levels of 638 are determined in this study using the MARVEL (Measured Active Rotational-Vibrational Energy Levels) procedure [6–9], built upon the theory of spectroscopic networks [10, 11], and the related code MARVEL 4.0.

The importance of the understanding of the energy-level structure and the high-resolution rovibrational spectra of CO_2 has been emphasized a number of times [1–4, 12, 13]. For example, carbon dioxide (a) is one of the most important constituents of a

This is an open access article under the terms of the [Creative Commons Attribution](https://creativecommons.org/licenses/by/4.0/) License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2024 The Author(s). *Journal of Computational Chemistry* published by Wiley Periodicals LLC.

TABLE 1 | Experimental sources of rovibrational transitions of $^{16}\text{O}^{13}\text{C}^{18}\text{O}$ used during the MARVEL analysis of this study, and certain characteristics of these sources.

Source	Range/ cm^{-1}	$A/V/D^a$	CSU^b	MSU^c
85Jolma [21]	578.71–722.69	335/335/0	5.0×10^{-3}	5.0×10^{-3}
98DeBeSmRi [22]	1211.71–1375.84	186/186/0	2.0×10^{-4}	3.1×10^{-4}
85Toth [23]	1224.55–1369.31	46/46/0	5.0×10^{-3}	5.0×10^{-3}
84RiBeDeFe [24]	1980.52–2614.51	39/39/0	1.0×10^{-3}	1.1×10^{-3}
85RiBeDe [25]	1980.52–2006.68	20/20/0	1.0×10^{-2}	1.0×10^{-2}
86EsSaRoVa [26]	2154.83–21812.84	796/796/0	9.0×10^{-4}	1.0×10^{-3}
83EsRo [27]	2157.47–2311.15	181/181/0	6.9×10^{-4}	8.1×10^{-4}
82EsHuSaVa [28]	2162.66–2311.15	176/176/0	1.0×10^{-3}	1.1×10^{-3}
86EsRo [29]	2165.05–2286.94	187/187/0	5.0×10^{-4}	9.8×10^{-4}
86BaRo [30]	2192.01–2305.11	163/163/0	1.0×10^{-3}	1.0×10^{-3}
12LyKaJaLu [31]	2194.32–4838.49	500/500/0	1.0×10^{-3}	1.3×10^{-3}
78BaLiDeRa [32]	2209.88–2298.64	86/86/0	3.0×10^{-3}	3.3×10^{-3}
68ObRaHaMc_PENN-STATE [33]	2235.89–2287.09	69/61/8	2.8×10^{-2}	3.1×10^{-2}
68ObRaHaMc_OHIO-STATE [33]	2235.92–2284.24	39/34/5	2.5×10^{-2}	3.5×10^{-2}
67Hahn [34]	2235.97–2273.03	39/34/5	1.0×10^{-3}	7.5×10^{-3}
03DeBeSmRi [35]	2240.56–2289.27	48/48/0	2.0×10^{-4}	2.4×10^{-4}
84DeRiBe [36]	2567.88–2614.51	27/27/0	1.0×10^{-3}	1.3×10^{-3}
13RoBoAuDr [37]	2682.92–2725.66	56/56/0	7.2×10^{-2}	7.2×10^{-2}
14BoJaLyTa [38]	3421.04–4679.99	398/398/0	3.0×10^{-4}	4.9×10^{-4}
08ToMiBrDe [39]	3437.73–3621.29	159/159/0	1.0×10^{-3}	1.1×10^{-3}
82BaRiSmRa [40]	3453.51–3607.95	79/79/0	5.0×10^{-3}	6.2×10^{-3}
03DiPeTaTe [41]	4449.06–8024.48	1372/1372/0	1.0×10^{-3}	1.5×10^{-3}
02Miller [42]	4469.44–4954.20	352/351/1	2.0×10^{-5}	1.2×10^{-4}
15BoJaLyTa [43]	4682.34–4946.90	103/103/0	4.7×10^{-4}	8.5×10^{-4}
18KaSiCeMo [44]	5698.72–5851.16	555/555/0	1.0×10^{-3}	1.1×10^{-3}
18CeKaMoKa [45]	5703.11–5832.29	158/158/0	1.0×10^{-3}	1.1×10^{-3}
18KaCeMoKa [46]	5703.11–5877.33	513/513/0	1.0×10^{-3}	1.1×10^{-3}
08PePeCa [47]	5851.97–7035.68	534/534/0	1.0×10^{-3}	1.3×10^{-3}
14KaCaMoKa [48]	5867.62–6744.16	1375/1375/0	1.0×10^{-3}	1.2×10^{-3}
06PeKaRoPe [49]	5957.24–6122.25	628/628/0	1.0×10^{-3}	1.3×10^{-3}
07PeKaRoPe [50]	5967.50–6319.32	41/41/0	1.0×10^{-3}	1.0×10^{-3}
04DiMaRoPe [51]	6099.92–6744.16	1313/1312/1	1.0×10^{-3}	1.6×10^{-3}
17KaCaKaTa [52]	6982.58–7917.50	516/515/1	1.0×10^{-3}	1.2×10^{-3}
10CaSoMoPe [53]	7029.57–7916.90	1044/1044/0	8.0×10^{-4}	9.4×10^{-4}
14KaKaTaPe [54]	7914.36–8024.24	92/92/0	1.5×10^{-3}	1.6×10^{-3}
05DiCaBeTa [55]	9153.61–9317.09	123/123/0	2.0×10^{-3}	3.5×10^{-3}

Note: ^a $A/V/D$ = Available/Validated/Deleted transitions (transitions in floating components cannot be validated, but are not deleted). ^b CSU = Average claimed source uncertainty. ^c MSU = Average MARVEL suggested uncertainty.

large number of planetary atmospheres, including that of Earth [14], where it has a significant contribution to the man-made greenhouse effect, (b) has an especially significant contribution to the radiative balance of the atmospheres of our neighboring planets Mars and Venus, and (c) has been used to establish the ratio of carbon isotopologues in the atmospheres of exoplanets [15]. Research on the high-resolution spectroscopy of carbon-dioxide isotopologues is very important as it supports such studies.

As to the energy-level structure of carbon dioxide, an especially restrictive rule applies for the symmetric CO_2 isotopologues: half of the rotational states are forbidden by the Pauli principle. Relaxation of this rule in asymmetric isotopologues increases the importance of these trace species, as all rotational states are allowed, leading to much denser spectra with lines displaced from those of the parent isotopologue. Thus, the many more allowed rovibrational transitions of 638 make their relative contribution to the detection of this molecule significantly more




FIGURE 1 | The number of transitions incident to each energy level versus the empirical rovibrational energy level values determined in this study.




FIGURE 2 | Rotational quantum number J for each calculated energy level versus the empirical energy levels determined for $^{16}\text{O}^{13}\text{C}^{18}\text{O}$ using the MARVEL procedure.

important than they would be based on the relative abundance of this minor isotopologue.

2 | Methodological Details

2.1 | Marvel

The MARVEL procedure [6–9] includes the construction of a spectroscopic network (SN) [10] from observed spectral line center positions, whereby each energy level (quantum state) serves as a vertex of the SN, and pairs of vertices are connected with the observed transitions forming the edges of the network. Under ideal circumstances, the larger the number of measured and collected spectral lines, the more empirical energy levels can be determined, or at least they are determined with higher accuracy. However, due to the limited number of experimental data available, one usually arrives at a fragmented SN, resulting in a principal component, where all the vertices are linked to the ground state, and a number of isolated components, which need to be connected to the principal one using accurately-known calculated or semi-empirical lines. For each spectral line, uncertainties, as well as unique labels, containing quantum numbers characterizing the upper and lower states, must be included in the input file for MARVEL. After executing the MARVEL procedure, the output file contains empirical energy levels, supplemented with educated estimates of their uncertainties. Here the bootstrap method implemented in MARVEL 4.0 [16] is used to determine the final uncertainties of the energy levels.

Inconsistencies, that is, line center positions that significantly deviate from the majority of the data provided are detected straightforwardly during a MARVEL run. This feature proves invaluable for identifying issues with the experimental data, whether stemming from user mistakes during data collection and analysis or from misassignments of the experimental features. The MARVEL input file was regularly tested not only for incorrectly labelled transitions, but also for those which do not obey well-established selection rules.

2.2 | Notation and Quantum Numbers

In its ground electronic state, CO₂ is a linear molecule; it has three fundamental vibrational modes, usually denoted as ν_1 (symmetric stretch mode), ν_2 (bend mode), and ν_3 (antisymmetric stretch mode), associated with the vibrational quantum numbers ν_i , $i = 1, 2$, and 3 , respectively. The doubly-degenerate ν_2 bending mode is characterized by an angular momentum, traditionally described by the quantum number ℓ . Due to Fermi-resonance interactions between the states $(\nu_1, (\nu_2 + 2)^\ell, \nu_3)$ and $(\nu_1 + 1, \nu_2^\ell, \nu_3)$, the vibrational states of CO₂ are customarily denoted by the so-called AFGL (Air Force Geophysics Laboratory) notation [17–19]. Using the AFGL notation, the vibrational energy levels are labeled as $(\nu_1, \nu_2, \ell, \nu_3, r)$, where r is the ranking index for states in Fermi resonance. The polyad number P , defined as $P = 2\nu_1 + \nu_2 + 3\nu_3$, collects the interacting states in blocks. P is not a quantum number, but behaves like one. None of these quantum numbers are “good” ones; thus, there are no strict selection rules for them.

For the purposes of a MARVEL analysis, each rovibrational state of CO₂ must be uniquely characterized by a set of descriptors. For 638, the set of descriptors used is $(J, \nu_1, \nu_2, \ell, \nu_3, r, e/f)$. The only good quantum number among the set of descriptors is J , associated with the rotational angular momentum. Allowed rovibrational transitions are governed by the selection rules $\Delta J = \pm 1$ and $\Delta J = 0$. Parity is also considered to be exact. Thus, the final descriptor we attach to a rovibrational energy level is the rotationless parity p , which is denoted here as either ‘e’ or ‘f’ [20]. The dipole selection rule for this parity is as follows: if $\Delta J = 0$, then $e \leftrightarrow f$, while if $\Delta J = \pm 1$, then $e \leftrightarrow e$ or $f \leftrightarrow f$. The upper and lower states involved in a transition are denoted by ν and ν' , respectively. The P, R, and Q transitions are specified using the lower-state rotational quantum number, J'' .

2.3 | Experimental Studies of Line Positions

Many rovibrational transitions for the 638 isotopologue of carbon dioxide have been detected experimentally [21–56]. The the

TABLE 2 | Experimentally determined vibrational band origins of the ¹⁶O¹³C¹⁸O molecule obtained in this study (‘MARVEL’) and their comparison with the corresponding entries of the CDS-296 (‘CDS’) dataset [12].

Label	MARVEL energy/cm ⁻¹	CDS energy/cm ⁻¹
0 0 0 0 1 e	0.0000 (0)	0.000000
0 1 0 0 2 e	1244.8997 (5)	1244.899 743
0 0 0 0 1 1 e	2265.9709 (11)	2265.971 552
0 2 0 0 1 e	2701.9561 (740)	2701.936 293
0 1 0 0 1 2 e	3490.3964 (5)	3490.394 853
0 1 0 0 1 1 e	3587.5468 (5)	3587.548 613
0 0 0 0 2 1 e	4508.7466 (11)	4508.747 345
0 2 0 0 1 3 e	4692.1737 (5)	4692.178 083
0 2 0 0 1 2 e	4814.5601 (5)	4814.561 614
0 2 0 0 1 1 e	4925.0200 (5)	4925.017 861
0 1 0 0 2 2 e	5712.6091 (1)	5712.609 317
0 1 0 0 2 1 e	5809.8644 (30)	5809.861 989
0 3 0 0 1 4 e	5876.5947 (11)	5876.595 711
0 3 0 0 1 3 e	6026.6241 (11)	6026.624 885
0 3 0 0 1 2 e	6140.1225 (11)	6140.122 650
0 3 0 0 1 1 e	6279.4881 (11)	6279.488 663
0 0 0 0 3 1 e	6728.3543 (11)	6728.354 758
0 2 0 0 2 2 e	7017.8905 (11)	7017.891 780
0 4 0 0 1 4 e	7220.7354 (9)	7220.737 827
0 4 0 0 1 3 e	7351.69067 (9)	7351.691 716
0 1 0 0 3 2 e	7911.5688 (9)	7911.569 220
0 2 0 0 3 2 e	9198.1886 (20)	9198.189 034




FIGURE 3 | The polyad number, $P = 2v_1 + v_2 + 3v_3$, for each empirical energy level versus the energy levels calculated using MARVEL.




FIGURE 4 | The ratio between the transition uncertainties suggested by MARVEL to the transition uncertainties given by the source articles versus the transition wavenumbers.

experimental source of rovibrational transitions collected are summarized in Table 1. In total, 12362 experimentally measured transitions have been collected from 35 literature sources, all of them are within the range $578 - 9318 \text{ cm}^{-1}$. Of all the experimentally measured transitions, only 7432 are unique. During the MARVEL analysis, 21 transitions had to be removed from the spectroscopic network (SN) set up. They are listed with negative wavenumber entries in the transition file given in the [Supplementary Information](#) to this paper.

The SN of the experimentally measured transitions of 638 contains four large floating components. These floating components contain 48 rovibrational energy levels (vertices). To connect these floating components to the principal component, we used calculated transitions taken from the Carbon Dioxide Spectroscopic Databank (CDSD-296) [12]. Altogether 14 CDSD-296 transitions, with $5 \times 10^{-3} \text{ cm}^{-1}$ uncertainty, were added to the MARVEL database of experimentally measured transitions. The tag of these lines in the input transition file is '19CDSD'.




FIGURE 5 | Fermi number r for each calculated energy level versus the empirical rovibrational energy levels determined in this study using MARVEL.




FIGURE 6 | Comparison between rovibrational energies of the present $^{16}\text{O}^{13}\text{C}^{18}\text{O}$ dataset and those of CDS-296 [12] (blue squares) and Ames-2021 [13] (red circles).

3 | Results and Discussion

3.1 | Main Results

Using the MARVEL procedure, 3975 empirical rovibrational energy levels have been obtained for 638, determined by 12 348/7432 measured/unique observed transitions. Figure 1 shows the number of transitions incident to each energy level and their distribution as a function of energy. This figure confirms that the degree distribution of the nodes of spectroscopic networks formed by experimentally measured transitions appear to be heavy-tailed [57]. The consequences of are as

follows: (a) the appearance of “hubs,” that is, a small number of highly interconnected nodes, (b) a connection preference that is, generally disassortative [58], that is the SN’s high-degree vertices preferentially attach to low-degree ones, (c) considerable robustness and error tolerance, and (d) an “ultra-small-world” property, that is the average length of the shortest paths scales as $\sim \log \log N$, where N is the number of nodes in the SN.

The empirical rovibrational energy levels determined in this study have J values up to 82, as seen in Figure 2. Each dotted curve in Figure 2 corresponds to a vibrational band, with

TABLE 3 | The set of empirical ('MARVEL') energy levels of this study with differences larger than 0.05cm^{-1} compared to their CDS-296 ('CDS-296') [12] counterparts.

Labels	MARVEL energy/ cm^{-1}	CDS-296 Energy/ cm^{-1}	Source
27 5 1 1 0 6 f	6915.7291	6915.9096	07PeKaRoPe [50]
28 5 1 1 0 6 f	6936.5410	6936.7477	07PeKaRoPe [50]
29 3 1 1 1 3 f	6957.9903	6958.0590	06PeKaRoPe [49]
29 5 1 1 0 6 f	6958.2380	6958.3287	07PeKaRoPe [50]
30 5 1 1 0 6 f	6980.5175	6980.6523	07PeKaRoPe [50]
31 5 1 1 0 6 e	7001.1083	7001.3042	07PeKaRoPe [50]
32 5 1 1 0 6 e	7024.6895	7024.9583	07PeKaRoPe [50]
32 3 1 1 1 3 e	7024.9128	7024.8047	07PeKaRoPe [50]
33 5 1 1 0 6 e	7049.2272	7049.3500	07PeKaRoPe [50]
57 3 0 0 1 2 e	7344.3153	7344.3926	04DiMaRoPe [51]
35 4 0 0 1 4 e	7681.4510	7681.3943	10CaSoMoPe [53]
36 4 0 0 1 4 e	7707.6009	7707.7004	10CaSoMoPe [53]
36 6 0 0 0 7 e	7707.8968	7707.9952	10CaSoMoPe [53]

the dots representing different J values. The energies of the 22 vibrational band origins (VBO), where $J=0$, are listed in Table 2. The stacks of energy levels clearly show the semirigid character of the CO_2 molecule. Note also the nice agreement between the MARVEL-based VBOs determined in this study and those reported in the CDS-296 databank [12].

Figure 3 shows the polyad numbers, P , for the empirical energy levels of 638 as a function of energy. The values range from 0 up to 13. Note also the nearly linear, stepwise increase of P with the increase of the energy.

Figure 4 shows the ratio between the transition uncertainties suggested by MARVEL and the transition uncertainties given by the source of the transition. Our analysis revealed that the uncertainties of about 170 transitions should be increased by one order of magnitude compared to the uncertainties declared in their sources, while the uncertainties of three transitions should be increased by two orders of magnitude. A comparison between the average claimed source uncertainty (CSU) and the average MARVEL suggested uncertainty (MSU) can be found

in Table 1. This table shows that the two values, CSU and MSU , have very good overall agreement, with the exception of one source, 02Miller [42], where our analysis suggests a value one order of magnitude higher compared to the uncertainty suggested by the source.

Figure 5 gives an idea about Fermi number r for each calculated energy level, where just 10, 7, and 1 energy levels have Fermi resonance numbers 5, 6, and 7, respectively.

3.2 | Comparison with Other Databases

All the empirical rovibrational energy levels reported in this study were compared with energy levels available in the databases Carbon Dioxide Spectroscopic Databank (CDS-296) [12] and NASA Ames-2021 [13]. These comparisons, see Figure 6, show good overall agreement.

As seen in Figure 6, our data are in significantly better agreement with CDS-296, with an average deviation of only 0.002cm^{-1} , than with Ames-2021, where the average deviation is 0.024cm^{-1} . This is not surprising, as the CDS-296 data are semiempirical in nature. Figure 6 shows that there are some MARVEL energy levels with deviations from the CDS-296 results that are significantly larger than the average. We collected these energy levels, involving the sources 04DiMaRoPe [51], 07PeKaRoPe [50], and 10CaSoMoPe [53], in Table 3. Since each energy level is determined by only one source, we could not decide whether the MARVEL or the CDS-296 results were correct. Settling these issues requires additional measurements of transitions involving these large- J quantum states.

4 | Summary and Conclusions

We presented a comprehensive analysis of all the available high-resolution, rovibrational transitions of the fifth most abundant isotopologue of carbon dioxide, $^{16}\text{O}^{13}\text{C}^{18}\text{O}$. For this computational analysis, the MARVEL algorithm and code [6–8] were employed. The transitions forming the basis of the MARVEL analysis were collected from 35 literature sources. The 12362 measured transitions collected for $^{16}\text{O}^{13}\text{C}^{18}\text{O}$ cover the wavenumber range of $578 - 9318\text{cm}^{-1}$, with the polyad number, $P = 2\nu_1 + \nu_2 + 3\nu_3$, ranging from 1 to 13. The number of unique transitions this database contains is 7432. Just 21 of all the transitions collected had to be excluded from our analysis.

The 12362 measured transitions determine 3975 empirical energy levels, extending up to 9800cm^{-1} . The average uncertainty of the levels is 0.0025cm^{-1} . Detailed comparisons with the CDS-296 [12] and the NASA Ames-2021 [13] databases reveal average differences, with respect to this work, of 0.002 and 0.024cm^{-1} , respectively. Note that CO_2 transition intensities can be computed very accurately using *ab initio* theory [13, 59, 60]; these can be combined with MARVEL energy levels, derived here, to provide highly accurate line lists [61], which can be difficult to obtain experimentally for trace species like $^{16}\text{O}^{13}\text{C}^{18}\text{O}$.

Acknowledgments

The authors thank STFC for funding the UK – Jordan collaboration under the Newton Fund grant ST/T001429/1. JT acknowledges the support of the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation program through Advance Grant number 883830. The work in Budapest has received funding from the HUN-REN Hungarian Research Network and the National Research, Development and Innovation Office (NKFIH, grant no. K138233). This publication supports research performed within the COST Action CA21101 “Confined molecular systems: from a new generation of materials to the stars” (COSY), funded by the European Cooperation in Science and Technology (COST). Ala'a Azzam thanks AstroJo Institute for providing the necessary support to complete this research project.

Data Availability Statement

The data involved in this paper is given in the [supporting information](#) to this paper.

References

- I. E. Gordon, L. S. Rothman, R. J. Hargreaves, et al., “The HITRAN2020 Molecular Spectroscopic Database,” *Journal of Quantitative Spectroscopy and Radiation Transfer* 277 (2022): 107949, <https://doi.org/10.1016/j.jqsrt.2021.107949>.
- M. T. I. Ibrahim, D. Alatoom, T. Furtenbacher, et al., “MARVEL Analysis of High-Resolution Rovibrational Spectra of $^{16}\text{O}^{13}\text{C}^{18}\text{O}$,” *Journal of Computational Chemistry* 45 (2024): 969–984, <https://doi.org/10.1002/jcc.27266>.
- D. Alatoom, M. T. I. Ibrahim, T. Furtenbacher, et al., “MARVEL Analysis of High-Resolution Rovibrational Spectra of $^{16}\text{O}^{12}\text{C}^{18}\text{O}$,” *Journal of Computational Chemistry* 45 (2024): 2558–2573, <https://doi.org/10.1002/jcc.27453>.
- A. A. A. Azzam, S. A. A. Azzam, K. A. A. Aburummana, et al., “MARVEL Analysis of High-Resolution Rovibrational Spectra of the $^{18}\text{O}^{12}\text{C}^{18}\text{O}$, $^{17}\text{O}^{12}\text{C}^{18}\text{O}$, and $^{18}\text{O}^{13}\text{C}^{18}\text{O}$ Isotopologues of Carbon Dioxide,” *Journal of Molecular Spectroscopy* 405 (2024): 111947.
- A. A. A. Azzam, B. M. J. A. Doud, M. Q. A. Shersheer, et al., “MARVEL Analysis of High-Resolution Rovibrational Spectra of $^{16}\text{O}^{12}\text{C}^{16}\text{O}$,” *Scientific Data* (2024).
- A. G. Császár, G. Czakó, T. Furtenbacher, and E. Mátyus, “An Active Database Approach to Complete Rotational–Vibrational Spectra of Small Molecules,” *Annual Reports in Computational Chemistry* 3 (2007): 155–176, [https://doi.org/10.1016/S1574-1400\(07\)03009-5](https://doi.org/10.1016/S1574-1400(07)03009-5).
- T. Furtenbacher, A. G. Császár, and J. Tennyson, “MARVEL: Measured Active Rotational–Vibrational Energy Levels,” *Journal of Molecular Spectroscopy* 245 (2007): 115–125, <https://doi.org/10.1016/j.jjms.2007.07.005>.
- T. Furtenbacher and A. G. Császár, “MARVEL: Measured Active Rotational–Vibrational Energy Levels. II. Algorithmic Improvements,” *Journal of Quantitative Spectroscopy and Radiation Transfer* 113 (2012): 929–935, <https://doi.org/10.1016/j.jqsrt.2012.01.005>.
- P. Árendás, T. Furtenbacher, and A. G. Császár, “On Spectra of Spectra,” *Journal of Mathematical Chemistry* 54 (2016): 806–822, <https://doi.org/10.1007/s10910-016-0591-1>.
- A. G. Császár and T. Furtenbacher, “Spectroscopic Networks,” *Journal of Molecular Spectroscopy* 266 (2011): 99–103, <https://doi.org/10.1016/j.jms.2011.03.031>.
- T. Furtenbacher and A. G. Császár, “The Role of Intensities in Determining Characteristics of Spectroscopic Networks,” *Journal of*

Molecular Structure 1009 (2012): 123–129, <https://doi.org/10.1016/j.molstruc.2011.10.057>.

- S. A. Tashkun, V. I. Perevalov, R. R. Gamache, and J. Lamouroux, “CDSD-296, High-Resolution Carbon Dioxide Spectroscopic Database: An Update,” *Journal of Quantitative Spectroscopy and Radiation Transfer* 228 (2019): 124–131, <https://doi.org/10.1016/j.jqsrt.2019.03.001>.
- X. Huang, D. W. Schwenke, R. S. Freedman, and T. J. Lee, “Ames-2021 CO_2 Dipole Moment Surface and IR Line Lists: Toward 0.1% Uncertainty for CO_2 IR Intensities,” *Journal of Physical Chemistry. A* 126 (2022): 5940–5964, <https://doi.org/10.1021/acs.jpca.2c01291>.
- K. P. Shine and G. E. Perry, “Radiative Forcing due to Carbon Dioxide Decomposed Into Its Component Vibrational Bands,” *Quarterly Journal of the Royal Meteorological Society* 149, no. 754 (2023): 1856–1866, <https://doi.org/10.1002/qj.4485>.
- Y. Zhang, I. A. G. Snellen, A. J. Bohn, et al., “The ^{13}CO -Rich Atmosphere of a Young Accreting Super-Jupiter,” *Nature* 595 (2021): 370–372, <https://doi.org/10.1038/s41586-021-03616-x>.
- J. Tennyson, T. Furtenbacher, S. N. Yurchenko, and A. G. Császár, “Empirical Rovibrational Energy Levels for Nitrous Oxide,” *Journal of Quantitative Spectroscopy and Radiation Transfer* 316 (2024): 108902, <https://doi.org/10.1016/j.jqsrt.2024.108902>.
- G. Amat and M. Pimbert, “On Fermi Resonance in Carbon Dioxide,” *Journal of Molecular Spectroscopy* 16 (1965): 278–290, [https://doi.org/10.1016/0022-2852\(65\)90123-2](https://doi.org/10.1016/0022-2852(65)90123-2).
- L. S. Rothman and L. D. G. Young, “Infrared Energy Levels and Intensities of Carbon Dioxide-II,” *Journal of Quantitative Spectroscopy and Radiation Transfer* 25 (1981): 505–524, [https://doi.org/10.1016/0022-4073\(81\)90026-1](https://doi.org/10.1016/0022-4073(81)90026-1).
- R. A. Toth, L. R. Brown, C. E. Miller, V. M. Devi, and D. C. Benner, “Spectroscopic Database of CO_2 Line Parameters: 4300–7000 cm^{-1} ,” *Journal of Quantitative Spectroscopy and Radiation Transfer* 109 (2008): 906–921, <https://doi.org/10.1016/j.jqsrt.2007.12.004>.
- J. M. Brown, J. T. Hougen, K. P. Huber, et al., “Labeling of Parity Doublet Levels in Linear Molecules,” *Journal of Molecular Spectroscopy* 55 (1975): 500–503, [https://doi.org/10.1016/0022-2852\(75\)90291-X](https://doi.org/10.1016/0022-2852(75)90291-X).
- K. Jolma, “Infrared Spectrum of Isotopic Carbon Dioxide in the Region of Bending Fundamental ν_2 ,” *Journal of Molecular Spectroscopy* 111 (1985): 211–218, [https://doi.org/10.1016/0022-2852\(85\)90002-5](https://doi.org/10.1016/0022-2852(85)90002-5).
- V. M. Devi, D. C. Benner, M. A. H. Smith, and C. P. Rinsland, “Pressure Broadening and Pressure Shift Coefficients in the $2\nu_2^0$ and ν_1 Bands of $^{16}\text{O}^{13}\text{C}^{18}\text{O}$,” *Journal of Quantitative Spectroscopy and Radiation Transfer* 60 (1998): 771–783, [https://doi.org/10.1016/S0022-4073\(98\)00081-8](https://doi.org/10.1016/S0022-4073(98)00081-8).
- R. A. Toth, “Line Positions and Strengths of CO_2 in the 1200–1430- cm^{-1} Region,” *Applied Optics* 24 (1985): 261–274, <https://doi.org/10.1364/AO.24.000261>.
- C. P. Rinsland, D. C. Benner, V. M. Devi, P. S. Ferry, C. H. Sutton, and D. J. Richardson, “Atlas of High Resolution Infrared Spectra of Carbon Dioxide,” *Applied Optics* 23 (1984): 2051–2052, <https://doi.org/10.1364/AO.23.002051>.
- C. P. Rinsland, D. C. Benner, and V. M. Devi, “Measurements of Absolute Line Intensities in Carbon Dioxide Bands Near 5.2 μm ,” *Applied Optics* 24 (1985): 1644–1650, <https://doi.org/10.1364/AO.24.001644>.
- M. P. Esplin, H. Sakai, L. S. Rothman, G. A. Vanasse, W. M. Barowy, and R. J. Huppi, *Carbon Dioxide Line Positions in the 2.8 and 4.3 Micron Regions at 800 Kelvin*, Tech. Rep. AFGL-TR-86-0046, (Logan, UT: Utah State University, 1986), <https://apps.dtic.mil/sti/citations/ADA173808>.
- M. P. Esplin and L. S. Rothman, “Spectral Measurements of High Temperature Isotopic Carbon Dioxide in the 4.3 μm Region,” *Journal of Molecular Spectroscopy* 100 (1983): 193–204, [https://doi.org/10.1016/0022-2852\(83\)90036-X](https://doi.org/10.1016/0022-2852(83)90036-X).

28. M. P. Esplin, R. J. Huppi, H. Sakai, G. A. Vanasse, and L. S. Rothman, *Absorption Measurements of CO₂ and H₂O at High Resolution and Elevated Temperatures*, Tech. Rep. AFGL-TR-82-0057 (Logan, UT: Utah State University, 1982), <https://apps.dtic.mil/sti/citations/ADA113824>.
29. M. P. Esplin and L. S. Rothman, "Spectral Measurements of High-Temperature Isotopic Carbon Dioxide in the 4.5- and 2.8 μm Regions," *Journal of Molecular Spectroscopy* 116 (1986): 351–363, [https://doi.org/10.1016/0022-2852\(86\)90132-3](https://doi.org/10.1016/0022-2852(86)90132-3).
30. D. Bailly and C. Rossetti, "¹³C¹⁶O¹⁸O: $\Sigma - \Sigma$ and $\Pi - \Pi$ Transitions in CO₂;N₂;O₂;He DC Discharges From High Information Emission Fourier Spectra," *Journal of Molecular Spectroscopy* 119 (1986): 388–397, [https://doi.org/10.1016/0022-2852\(86\)90033-0](https://doi.org/10.1016/0022-2852(86)90033-0).
31. O. M. Lyulin, E. V. Karlovets, D. Jacquemart, Y. Lu, A. W. Liu, and V. I. Perevalov, "Infrared Spectroscopy of ¹⁷O- and ¹⁸O-Enriched Carbon Dioxide in the 1700–8300 cm⁻¹ Wavenumber Region," *Journal of Quantitative Spectroscopy and Radiation Transfer* 113 (2012): 2167–2181, <https://doi.org/10.1016/j.jqsrt.2012.06.028>.
32. A. Baldacci, L. Linden, V. M. Devi, K. N. Rao, and B. Fridovich, "Interpretation of the ¹³C¹⁶O₂ Spectrum at 4.4 μm ," *Journal of Molecular Spectroscopy* 72 (1978): 135–142, [https://doi.org/10.1016/0022-2852\(78\)90049-8](https://doi.org/10.1016/0022-2852(78)90049-8).
33. R. Oberly, K. N. Rao, Y. H. Hahn, and T. K. McCubbin, "Bands of Carbon Dioxide in the Region of 4.3 Microns," *Journal of Molecular Spectroscopy* 25 (1968): 138–165, [https://doi.org/10.1016/0022-2852\(68\)80002-5](https://doi.org/10.1016/0022-2852(68)80002-5).
34. Y. H. Hahn, "The Absorption and Emission Spectra of Carbon-Dioxide at 4.3 Microns," (PhD diss., The Pennsylvania State University 1967), <https://www.proquest.com/openview/074bf2696a7b47d2793ca1d620a02193/1?pq-origsite=gscholar&cbl=18750&diss=y>.
35. V. M. Devi, D. C. Benner, M. A. H. Smith, and C. P. Rinsland, "Nitrogen Broadening and Shift Coefficients in the 4.2–4.5 Micron Bands of CO₂," *Journal of Quantitative Spectroscopy and Radiative Transfer* 76 (2003): 289–307.
36. V. M. Devi, C. P. Rinsland, and D. C. Benner, "Absolute Intensity Measurements of CO₂ Bands in the 2395–2680-cm⁻¹ Region," *Applied Optics* 23, no. 22 (1984): 4067, <https://doi.org/10.1364/AO.23.004067>.
37. S. Robert, Y. Borkov, J. Vander Auwera, et al., "Assignment and Rotational Analysis of New Absorption Bands of Carbon Dioxide Isotopologues in Venus Spectra," *Journal of Quantitative Spectroscopy and Radiation Transfer* 114 (2013): 29–41, <https://doi.org/10.1016/j.jqsrt.2012.08.023>.
38. Y. G. Borkov, D. Jacquemart, O. M. Lyulin, S. A. Tashkun, and V. I. Perevalov, "Infrared Spectroscopy of ¹⁷O- and ¹⁸O-Enriched Carbon Dioxide: Line Positions and Intensities in the 3200–4700 cm⁻¹ Region. Global Modeling of the Line Positions of ¹⁶O¹²C¹⁷O and ¹⁷O¹²C¹⁷O," *Journal of Quantitative Spectroscopy and Radiation Transfer* 137 (2014): 57–76, <https://doi.org/10.1016/j.jqsrt.2013.11.008>.
39. R. Toth, C. Miller, L. Brown, V. M. Devi, and D. C. Benner, "Line Strengths of ¹⁶O¹³C¹⁶O, ¹⁶O¹³C¹⁸O, ¹⁶O¹³C¹⁷O and ¹⁸O¹³C¹⁸O Between 2200 and 6800 Cm⁻¹," *Journal of Molecular Spectroscopy* 251 (2008): 64–89, <https://doi.org/10.1016/j.jms.2008.01.009>.
40. A. Baldacci, C. P. Rinsland, M. A. H. Smith, and K. N. Rao, "Spectrum of ¹³C¹⁶O₂ at 2.8 μm ," *Journal of Molecular Spectroscopy* 94 (1982): 351–362, [https://doi.org/10.1016/0022-2852\(82\)90011-X](https://doi.org/10.1016/0022-2852(82)90011-X).
41. Y. Ding, V. Perevalov, S. Tashkun, J.-L. Teffo, A.-W. Liu, and S.-M. Hu, "¹⁶O¹³C¹⁸O: High-Resolution Absorption Spectrum Between 4000 and 9500cm⁻¹ and Global Fitting of Vibration-Rotational Line Positions," *Journal of Molecular Spectroscopy* 222 (2003): 276–283, <https://doi.org/10.1016/j.jms.2003.08.004>.
42. C. E. Miller, "Private communication," 2024.
43. Y. G. Borkov, D. Jacquemart, O. M. Lyulin, S. A. Tashkun, and V. I. Perevalov, "Infrared Spectroscopy of ¹⁷O- and ¹⁸O-Enriched Carbon Dioxide: Line Positions and Intensities in the 4681–5337 cm⁻¹ Region," *Journal of Quantitative Spectroscopy and Radiation Transfer* 159 (2015): 1–10, <https://doi.org/10.1016/j.jqsrt.2015.02.019>.
44. E. V. Karlovets, A. D. Sidorenko, P. Čermák, et al., "The ¹³CO₂ Absorption Spectrum by CRDS Near 1.74 μm ," *Journal of Molecular Spectroscopy* 354 (2018): 54–59, <https://doi.org/10.1016/j.jms.2018.10.003>.
45. P. Čermák, E. V. Karlovets, D. Mondelain, S. Kassi, V. I. Perevalov, and A. Campargue, "High Sensitivity CRDS of CO₂ in the 1.74 μm Transparency Window. A Validation Test for the Spectroscopic Databases," *Journal of Quantitative Spectroscopy and Radiation Transfer* 207 (2018): 95–103, <https://doi.org/10.1016/j.jqsrt.2017.12.018>.
46. E. V. Karlovets, P. Čermák, D. Mondelain, et al., "Analysis and Theoretical Modeling of the ¹⁸O Enriched Carbon Dioxide Spectrum by CRDS Near 1.74 μm ," *Journal of Quantitative Spectroscopy and Radiation Transfer* 217 (2018): 73–85, <https://doi.org/10.1016/j.jqsrt.2018.05.017>.
47. B. V. Perevalov, V. I. Perevalov, and A. Campargue, "A (Nearly) Complete Experimental Linelist for ¹³C¹⁶O₂, ¹⁶O¹³C¹⁸O, ¹⁶O¹³C¹⁷O, ¹³C¹⁸O₂ and ¹⁷O¹³C¹⁸O by High-Sensitivity CW-CRDS Spectroscopy Between 5851 and 7045 cm⁻¹," *Journal of Quantitative Spectroscopy and Radiation Transfer* 109 (2008): 2437–2462, <https://doi.org/10.1016/j.jqsrt.2008.03.010>.
48. E. V. Karlovets, A. Campargue, D. Mondelain, S. Kassi, S. A. Tashkun, and V. I. Perevalov, "High Sensitivity Cavity Ring Down Spectroscopy of ¹⁸O Enriched Carbon Dioxide Between 5850 and 7000 cm⁻¹: Part II Analysis and Theoretical Modeling of the ¹²C¹⁸O₂, ¹³C¹⁸O₂ and ¹⁶O¹³C¹⁸O Spectra," *Journal of Quantitative Spectroscopy and Radiation Transfer* 136 (2014): 71–88, <https://doi.org/10.1016/j.jqsrt.2013.11.005>.
49. B. V. Perevalov, S. Kassi, D. Romanini, V. I. Perevalov, S. A. Tashkun, and A. Campargue, "CW-Cavity Ringdown Spectroscopy of Carbon Dioxide Isotopologues Near 1.5 μm ," *Journal of Molecular Spectroscopy* 238 (2006): 241–255, <https://doi.org/10.1016/j.jms.2006.05.009>.
50. B. V. Perevalov, S. Kassi, D. Romanini, V. I. Perevalov, S. A. Tashkun, and A. Campargue, "Global Effective Hamiltonians of ¹⁶O¹³C¹⁷O and ¹⁶O¹³C¹⁸O Improved From CW-CRDS Observations in the 5900–7000 cm⁻¹ Region," *Journal of Molecular Spectroscopy* 241 (2007): 90–100, <https://doi.org/10.1016/j.jms.2006.11.003>.
51. Y. Ding, P. Macko, D. Romanini, et al., "High Sensitivity CW-Cavity Ringdown and Fourier Transform Absorption Spectroscopies of ¹³CO₂," *Journal of Molecular Spectroscopy* 226 (2004): 146–160, <https://doi.org/10.1016/j.jms.2004.03.009>.
52. E. V. Karlovets, A. Campargue, S. Kassi, S. A. Tashkun, and V. I. Perevalov, "Analysis and Theoretical Modeling of ¹⁸O Enriched Carbon Dioxide Spectrum by CRDS Near 1.35 μm : (II) ¹⁶O¹³C¹⁸O, ¹⁶O¹³C¹⁷O, ¹²C¹⁸O₂, ¹⁷O¹²C¹⁸O, ¹²C¹⁷O₂, ¹³C¹⁸O₂ and ¹⁷O¹³C¹⁸O," *Journal of Quantitative Spectroscopy and Radiation Transfer* 191 (2017): 75–87, <https://doi.org/10.1016/j.jqsrt.2017.01.038>.
53. A. Campargue, K. F. Song, N. Mouton, V. I. Perevalov, and S. Kassi, "High Sensitivity CW-Cavity Ring Down Spectroscopy of Five ¹³CO₂ Isotopologues of Carbon Dioxide in the 1.26–1.44 μm Region (I): Line Positions," *Journal of Quantitative Spectroscopy and Radiation Transfer* 111 (2010): 659–674, <https://doi.org/10.1016/j.jqsrt.2009.11.013>.
54. E. V. Karlovets, S. Kassi, S. A. Tashkun, V. I. Perevalov, and A. Campargue, "High Sensitivity Cavity Ring Down Spectroscopy of Carbon Dioxide in the 1.19–1.26 μm Region," *Journal of Quantitative Spectroscopy and Radiation Transfer* 144 (2014): 137–153, <https://doi.org/10.1016/j.jqsrt.2014.04.001>.
55. Y. Ding, A. Campargue, E. Bertseva, S. Tashkun, and V. I. Perevalov, "Highly Sensitive Absorption Spectroscopy of Carbon Dioxide by ICLAS-VeCSEL Between 8800 and 9530 cm⁻¹," *Journal of Molecular Spectroscopy* 231 (2005): 117–123, <https://doi.org/10.1016/j.jms.2004.12.008>.

56. D. C. Benner, V. M. Devi, K. Sung, et al., "Line Parameters Including Temperature Dependences of Air- and Self-Broadened Line Shapes of $^{12}\text{C } ^{16}\text{O}_2$: 2.06- μm Region," *Journal of Molecular Spectroscopy* 326 (2016): 21–47, <https://doi.org/10.1016/j.jms.2016.02.012>.
57. T. Furtenbacher, P. Árendás, G. Mellau, and A. G. Császár, "Simple Molecules as Complex Systems," *Scientific Reports* 4 (2014): 4654, <https://doi.org/10.1038/srep04654>.
58. M. E. J. Newman, *Networks* (Oxford: Oxford University Press, 2010).
59. O. L. Polyansky, K. Bielska, M. Ghysels, et al., "High Accuracy CO_2 Line Intensities Determined From Theory and Experiment," *Physical Review Letters* 114 (2015): 243001, <https://doi.org/10.1103/PhysRevLett.114.243001>.
60. T. Odintsova, E. Fasci, L. Moretti, et al., "Highly-Accurate Intensity Factors of Pure CO_2 Lines Near 2 μm ," *Journal of Chemical Physics* 146 (2017): 244309, <https://doi.org/10.1063/1.4989925>.
61. E. J. Zak, J. Tennyson, O. L. Polyansky, et al., "Room Temperature Line Lists for CO_2 Asymmetric Isotopologues with *Ab Initio* Computed Intensities," *Journal of Quantitative Spectroscopy and Radiation Transfer* 203 (2017): 265–281, <https://doi.org/10.1016/j.jqsrt.2017.01.037>.

Supporting Information

Additional supporting information can be found online in the Supporting Information section.