

Refined line lists for the ions of ArH⁺, KrH⁺ and XeH⁺

Ben Such, Samuel Hamilton, Aberystwyth University; Maire N. Gorman,¹ Aberystwyth University / UCL; Sergey N. Yurchenko, Jonathan Tennyson, UCL

Here we present refined ground state ($X^1\Sigma^+$) line lists for the ions of ArH⁺, KrH⁺ and XeH⁺. The various isotopologues considered are ³⁶ArH⁺, ³⁸ArH⁺, ⁴⁰ArH⁺, ⁸⁴KrH⁺, ¹²⁹XeH⁺, ¹³¹XeH⁺ and ¹³²XeH⁺. To date, ³⁶ArH⁺ has been found in the Crab Nebula and both ³⁶ArH⁺ and ³⁸ArH⁺ have been detected in the direction of the lensed blazar PKS 1830 - 211. As far as we are aware neither KrH⁺ or XeH⁺ have been detected in space. *Ab initio* Potential Energy and Dipole Moment curves were calculated using MRCI/aug-cc-pVQZ level of theory and the former refined by fitting experimental data to an Extended Morse Oscillator to accuracies of ~ 0.60 cm⁻¹ (ArH⁺) and ~ 0.0009 cm⁻¹ (KrH⁺, XeH⁺). All line lists contain 400 states and 4000 - 7000 transitions covering appropriate vibrational and rotational states up to dissociation. Further work will now include adapting the Dipole moment curves to take into account the difference in position between centre of mass and centre of charge for these ions.

¹mng2@aber.ac.uk